



ISSN : 2347 - 2243

Indo - American Journal of Life Sciences and Biotechnology



<https://iajlb.org>

Email : editor@iajlb.com or iajlb.editor@gmail.com



Optimisation of a Rapid CTAB Protocol for High-Quality DNA Extraction From Polysaccharide-Rich Plant Tissues

Francesca Bellandi¹●, Garrett Whitfield²●, Eleanor Rowntree³●

¹Associate Professor, Department of Plant Molecular Biology, Sannio Institute of Pharmacology, Benevento, Italy.

²Assistant Professor, Department of Molecular Biology, Cumberland Plateau Institute of Life Sciences, Cookeville, USA.

³Assistant Professor, Department of Biotechnology, Penwith Institute of Applied Microbiology, Penzance, United Kingdom.

Email: ¹f.bellandi@sip-bn.it, ²g.whitfield@cpils.edu, ³e.rowntree@piam.ac.uk

ORCID: ¹0000-0001-1107-2972, ²0000-0001-0827-0839, ³0000-0001-1125-7248

Received: September 11, 2012 Accepted: December 20, 2012 Published: April 18, 2013

DOI: [10.62644/iajlb.2013.v10.i2.pp11-18](https://doi.org/10.62644/iajlb.2013.v10.i2.pp11-18)

ABSTRACT

The cetyltrimethylammonium bromide (CTAB) method has been the workhorse of plant DNA extraction for more than three decades, yet it is notorious for failing on tissues loaded with polysaccharides and polyphenols. Mucilage, gums, pectins and storage carbohydrates co-precipitate with nucleic acids, turn pellets into glassy gels, inhibit polymerases and restriction enzymes, and distort absorbance readings. This narrative review asks what the published literature says about making a CTAB protocol both rapid and reliable for such tissues. It draws on the original CTAB papers, on the many modifications proposed for difficult species, and on work about PCR inhibition and the assessment of DNA quality, and it reports no new laboratory data. Optimisation emerges as a matter of balance rather than of any single magic additive: salt concentration, polyvinylpyrrolidone, reducing agents, chloroform steps and precipitation conditions each solve one problem while risking another, and speed gained by shortening a step is often lost again in repeated extractions. Quality checks based on absorbance ratios are useful but easily misread, and amplification of the intended target remains the decisive test. The review closes with a short list of what a convincing optimisation report should contain.

Keywords: CTAB; DNA extraction; Polysaccharides; Polyphenols; PCR inhibition; Protocol optimisation; Narrative review

Citation: Bellandi, Whitfield and Rowntree [2013]. Optimisation of a Rapid CTAB Protocol for High-Quality DNA Extraction From Polysaccharide-Rich Plant Tissues. *Indo-Am. J. Life Sci. Biotechnol.* 10(2):11-18. DOI: [10.62644/iajlb.2013.v10.i2.pp11-18](https://doi.org/10.62644/iajlb.2013.v10.i2.pp11-18)

Article Class: Narrative Review

1. Introduction

Almost every plant molecular study begins with the same task: getting DNA out of a leaf. For many species it is easy; for those that store carbohydrate, secrete mucilage or accumulate phenolics, it is the step on

which the project stalls. The method most often reached for in these cases is extraction in cetyltrimethylammonium bromide, or CTAB, a detergent introduced for plant work in the early 1980s (1) and adapted for tiny samples soon afterwards



(2).

The appeal of CTAB is practical. It needs no specialised equipment, scales from a leaf disc to a large preparation, and under high-salt conditions it holds most polysaccharides in solution while nucleic acids are separated from them (1). Its reputation is nonetheless uneven: protocols that work for one crop give brown, viscous or unamplifiable DNA from another, and a large literature of modifications has grown up around the problem (3).

This article reviews what that literature says about making a CTAB protocol both rapid and dependable for polysaccharide-rich tissues. It is a narrative review and reports no laboratory data; although its title speaks of optimising a protocol, it describes no particular experiment, laboratory or species. The aim is a fair account of what is established and what is merely repeated from one protocol paper to the next.

2. Literature Review

2.1 Origins of the CTAB Method

Before CTAB, plant DNA was commonly prepared by adapting procedures developed for microorganisms, which relied on detergent lysis, deproteinisation with chloroform and alcohol precipitation (4). Plant tissue was a poorer fit, because walls resisted lysis and vacuolar contents attacked the DNA once released. The CTAB procedure used a cationic detergent that lyses membranes and complexes with nucleic acids, with a salt concentration chosen so that the complex stays soluble while most polysaccharides do not co-precipitate (1).

A scaled-down version handled milligram amounts of fresh tissue and even herbarium material (2). A parallel line of work used sodium dodecyl sulfate with potassium acetate (5), and rapid versions dispensed with phenol and chloroform

altogether (6). Those methods are fast, yet they tend to carry more carbohydrate and pigment into the final solution, which is exactly what difficult tissues punish.

2.2 What Makes Some Tissues Difficult

Three groups of compounds account for most of the trouble. Polysaccharides, such as pectins, gums, mucilage and storage carbohydrates, are structurally similar enough to nucleic acids to follow them through alcohol precipitation, where they give a glassy pellet that is hard to dissolve (7). Polyphenols oxidise when cells are broken, and the resulting quinones bind covalently to DNA, giving brown preparations that resist enzymes (8). Latex and resins complicate some families (9). Table 1 summarises the main contaminants, their effects and the usual countermeasures.

Table 1. Contaminants typical of polysaccharide-rich tissue and what they do to an extract.

Contaminant	Typical source	Effect	Usual counter-measure
Polysaccharides	Mucilage, pectin, starch	Viscous, glassy pellet; enzyme inhibition (10, 11)	High salt; extra chloroform steps (7)
Polyphenols	Tannins, oxidised phenolics	Brown, cross-linked DNA (8)	PVP, reducing agent (12, 13)
Latex and resins	Laticifers, resin ducts	Gummy lysate (9)	Washed tissue, extra extraction
Proteins and RNA	Cell lysate	Inflated A260 reading (14)	Chloroform; RNase treatment
Residual CTAB or salt	Carry-over from wash	Polymerase inhibition (15)	Ethanol wash, full drying

Note: Qualitative summary of effects described in the cited literature; severity varies by species and tissue.

Carbohydrate contamination is insidious: a heavily contaminated extract can read as acceptable at 260 nm, since



polysaccharides absorb little, while still inhibiting the enzymes that follow (10). Plant polysaccharides and polyphenols are common causes of failed reactions, and polymerases differ in how well they tolerate them (11, 15). Such differences look small at the bench, but they decide whether a downstream reaction runs at all.

2.3 Modifications for Polysaccharide-Rich Species

The best-known modifications shift the chemistry in one direction. Raising the salt concentration keeps more polysaccharide soluble and was a central element of protocols designed for cotton and grapevine, two crops that combine mucilage with phenolics (16, 17). Adding polyvinylpyrrolidone, usually with a reducing agent, binds phenolics before they oxidise (12, 13). One modification for plants rich in both polysaccharides and polyphenols combined several of these changes with extra extraction and washing (7). Work on latex-bearing species added extra extraction rounds (9), and a cotton protocol began by isolating nuclei (18).

Simplified versions have also appeared. A microprep for tomato and other herbaceous plants trades yield for speed (19), protocols exist that dispense with liquid nitrogen and phenol (20, 21), and a detailed protocol article describes rapid CTAB isolation from small tissue samples (22). Others go further, sending a crude lysate straight to amplification (23) or replacing manual handling with high-throughput formats (24). These work well on easy tissues and, as a rule, poorly on mucilaginous ones. This is why the same species can yield cleanly in one laboratory and badly in another.

2.4 Sample Handling Before Extraction

Much of what decides success happens before the buffer is added. Field tissue can be dried over silica gel (25) or kept in

saturated salt and CTAB solution (26). Herbarium specimens are harder, because storage fragments DNA, and protocols for them have been compared in some detail (27, 28).

3. Materials and Methods

3.1 Scope

The question is deliberately narrow: what does the published literature say about optimising a rapid CTAB protocol for polysaccharide-rich plant tissues? No particular species is assumed, and no data from any institution, laboratory or field site are reported; the title should not be read as describing an experiment by the authors.

3.2 Literature Assembly

This is a narrative synthesis. No formal systematic search or screening procedure was used. Sources were assembled from the authors' reading of foundational and methodological papers on plant DNA extraction, PCR inhibition and DNA quality published up to the end of 2012, plus reference chasing. Selection is therefore subject to the usual biases of a narrative review.

3.3 Appraisal Framework

Each source was read for what was changed, what was measured to show improvement and how far the result might transfer. Reports were treated as stronger when they tested more than one species, used a standard comparator and measured amplification or digestion as well as absorbance. Table 2 and Table 3 collect the buffer components and quality checks that recurred.

4. Results

4.1 The Typical Rapid CTAB Workflow

Across the sources, a rapid CTAB protocol follows a recognisable sequence, shown in Figure 1: grinding in warm buffer, a short incubation, extraction with chloroform and



isoamyl alcohol, alcohol precipitation, washing and dissolving. Each stage has a characteristic way of going wrong: phenolics oxidise at the start, mucilage emulsifies at the interface, the pellet turns gelatinous at precipitation, and an over-dried pellet resists dissolving. None of these steps is difficult, but each is easy to skip when the bench is busy.

Table 2. Components of a CTAB extraction buffer and the reason each is there.

Component	Role	Caution
CTAB	Detergent; lyses membranes, binds nucleic acids (1)	Precipitates at low salt
Sodium chloride	Keeps polysaccharides soluble (1, 7)	Too little leaves carbohydrate behind
Tris and EDTA	Buffer; chelates metals that help nucleases (2)	Weak buffering fails on acidic tissue
PVP	Binds polyphenols (12, 17)	Can carry over into the pellet
Reducing agent	Limits oxidation of phenolics (8, 13)	Odour; volatile; safety handling
Chloroform-isoamyl alcohol	Removes proteins and pigments (4)	Emulsions with mucilage

Note: Roles are those described in the cited protocols; amounts are deliberately omitted because they are tuned to the tissue.

What distinguishes a rapid protocol from a slow one is rarely the chemistry; it is how often each step is repeated and how long each incubation lasts. No general rule says how far these can be shortened before yield or purity suffers, and authors of rapid protocols seldom report failures, so the record probably favours successes. The point is not that a published protocol is poor, only that its success travels less far than its title suggests.




Figure 1. Flow of a CTAB extraction with the points at which polysaccharide trouble usually appears (schematic, not data).

4.2 Which Step Addresses Which Problem

The reported modifications can be arranged by the problem they address, and Figure 2 does this qualitatively. Salt concentration is the principal lever against polysaccharide; PVP and a reducing agent address polyphenol; chloroform extraction removes protein and pigment and also helps with some carbohydrate; and a careful wash is what removes carry-over of salt and detergent. The matrix summarises the authors' reading, not a measured result. A pellet that has been given time to dissolve behaves differently from one that has been rushed.

	Salt	PVP	Reducer	Solvent	Wash
Polysaccharide	main remedy			partial help	
Polyphenol		main remedy	main remedy	partial help	
Protein / RNA				main remedy	
Salt carry-over					main remedy

Figure 2. Qualitative matrix of contaminant problems and the steps that address them (schematic, not data).

No single component carries the weight. A protocol that raises salt but omits a reducing agent may handle mucilage and still give brown DNA from a phenolic tissue, and one that adds PVP but is



washed carelessly may inhibit the polymerase through residue (15, 29).

4.3 Assessing the Extract

Authors judge extracts by several means, listed in Table 3. Absorbance ratios are the commonest. A ratio of absorbance at 260 and 280 nm near 1.8 is conventionally taken as acceptable for DNA, and a ratio at 260 and 230 nm above about 2.0 as indicating freedom from salt and organic contaminants, but both depend on pH, ionic strength and the blank used (14, 30). Gel electrophoresis shows size and integrity and reveals RNA (31). The decisive tests are functional: a restriction digest, or amplification of the intended locus (32).

Table 3. Common quality checks for an extract and what each can and cannot tell.

Check	What it shows	Limitation
A260/A280	Protein or phenol carry-over (14, 30)	Insensitive to polysaccharide
A260/A230	Salt, carbohydrate, organic solvent (30)	Varies with buffer and blank
Gel electrophoresis	Size and integrity; RNA smear (31)	Not quantitative for purity
Restriction digest	Whether enzymes can cut (10)	Needs enzyme and substrate
Target PCR	Whether the intended locus amplifies (32)	Primer-specific; spike controls help

Note: Interpretation follows the cited methodological papers; no thresholds are proposed beyond the usual rules of thumb.

A recurring weakness is that reports of improved protocols rest mainly on absorbance and a gel, without amplification of more than one target and without a spike-in control that would separate inhibition from an absent template. Where such controls were used, an extract that looked pure by ratio could

still inhibit. Read this way, a failed amplification is a clue about the extract, not proof that the template is absent.

4.4 Downstream Demands

What counts as good enough depends on use. PCR (33, 34) tolerates some impurity and degradation, especially for short targets. Fingerprinting with arbitrary primers (35) or selective amplification (36) is more demanding, since a trace of inhibitor can change band patterns. Barcoding with plastid markers (37, 38, 39) and large-scale genotyping (40, 41) ask for DNA of consistent quality across many samples, a demand that sequencing platforms have sharpened (42).

5. Discussion

5.1 Interpretation

Read across these reports, optimisation of a rapid CTAB protocol is a matter of balance. Every remedy has a cost: more salt dissolves more carbohydrate but can leave salt behind, extra chloroform steps lower contamination but lose DNA at each interface, and PVP removes phenolics but can itself carry over. The protocols that work best on difficult tissue are rarely the shortest, and any that claim to be both fast and universal deserve scepticism.

A protocol developed on one species is, in effect, a hypothesis about another, since carbohydrate and phenolic profiles change with season, age and growing conditions. Figure 3 sketches how a poor extract might be read. A reader choosing a protocol needs the answer to a simple question: will this work on my tissue?

5.2 Limitations

This review has the limitations of its design. It is a narrative synthesis without a systematic search, so relevant reports may have been missed. Most of the protocol papers it draws on are short methods notes that rarely include negative results or



replicate comparisons, and comparisons across laboratories are confounded by differences in tissue, equipment and analyst skill. Reviews of this kind are more useful when they say where the evidence stops than when they smooth it over.




Figure 3. Decision tree for reading a poor extract (schematic, not data).

5.3 Future Work

A convincing optimisation report would state the tissue and its age, describe every step precisely enough to repeat, include a standard comparator, and report more than absorbance: at minimum a gel, a digest or target amplification with a spike-in control. It would give time, cost and failure rate, not only the best result. It is a habit worth keeping even when the extract looks fine.

6. Conclusion

CTAB remains a sound basis for extracting DNA from polysaccharide-rich plant tissue, but the literature shows that its success depends on matching the chemistry to the tissue rather than on any single additive. Salt, PVP, reducing agents, chloroform steps and washing each address a specific problem, and shortening the protocol must not remove the step that addresses the problem the tissue actually has. Quality should be judged by function, and a report of improvement should show amplification, not only absorbance ratios. That is a modest claim, and it is the one the literature will bear.

For the practitioner, the message is to treat a published protocol as a starting point, test it on the tissue in hand and record what is changed. For the researcher, the opportunity is a set of side-by-side comparisons that would turn anecdote into guidance. These points apply with equal force to any extraction chemistry, not only to CTAB.

7. Acknowledgements

The authors thank colleagues who commented on earlier drafts of this review.

8. Funding

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

9. Conflict of Interest

The authors declare no conflict of interest.

10. Author Contributions

F.B. conceived the review and wrote the sections on the CTAB method and its modifications. G.W. wrote the sections on contaminants and quality assessment. E.R. wrote the sections on sample handling and downstream use. All authors revised the manuscript and approved the final version.

11. Ethics Statement

This is a review of published literature. No human or animal subjects were involved and no ethical approval was required.

12. Data Availability

No new data were generated. All sources are cited in the reference list.

13. References

1. Murray, M. G., & Thompson, W. F. (1980). Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research*, 8(19), 4321–4326. DOI: [10.1093/nar/8.19.4321](https://doi.org/10.1093/nar/8.19.4321)
2. Rogers, S. O., & Bendich, A. J. (1985). Extraction of DNA from milligram amounts of fresh, herbarium and mummified plant tissues. *Plant Molecular Biology*, 5(2), 69–76. DOI:



- [10.1007/BF00020088](https://doi.org/10.1007/BF00020088)
3. Varma, A., Padh, H., & Shrivastava, N. (2007). Plant genomic DNA isolation: An art or a science. *Biotechnology Journal*, 2(3), 386–392. DOI: [10.1002/biot.200600195](https://doi.org/10.1002/biot.200600195)
 4. Marmur, J. (1961). A procedure for the isolation of deoxyribonucleic acid from micro-organisms. *Journal of Molecular Biology*, 3(2), 208–218. DOI: [10.1016/S0022-2836\(61\)80047-8](https://doi.org/10.1016/S0022-2836(61)80047-8)
 5. Dellaporta, S. L., Wood, J., & Hicks, J. B. (1983). A plant DNA miniprep: Version II. *Plant Molecular Biology Reporter*, 1(4), 19–21. DOI: [10.1007/BF02712670](https://doi.org/10.1007/BF02712670)
 6. Edwards, K., Johnstone, C., & Thompson, C. (1991). A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Research*, 19(6), 1349. DOI: [10.1093/nar/19.6.1349](https://doi.org/10.1093/nar/19.6.1349)
 7. Porebski, S., Bailey, L. G., & Baum, B. R. (1997). Modification of a CTAB DNA extraction protocol for plants containing high polysaccharide and polyphenol components. *Plant Molecular Biology Reporter*, 15(1), 8–15. DOI: [10.1007/BF02772108](https://doi.org/10.1007/BF02772108)
 8. Loomis, W. D. (1974). Overcoming problems of phenolics and quinones in the isolation of plant enzymes and organelles. *Methods in Enzymology*, 31, 528–544. DOI: [10.1016/S0076-6879\(74\)31057-9](https://doi.org/10.1016/S0076-6879(74)31057-9)
 9. Michiels, A., Van den Ende, W., Tucker, M., Van Riet, L., & Van Laere, A. (2003). Extraction of high-quality genomic DNA from latex-containing plants. *Analytical Biochemistry*, 315(1), 85–89. DOI: [10.1016/S0003-2697\(02\)00665-6](https://doi.org/10.1016/S0003-2697(02)00665-6)
 10. Wilson, I. G. (1997). Inhibition and facilitation of nucleic acid amplification. *Applied and Environmental Microbiology*, 63(10), 3741–3751. DOI: [10.1128/aem.63.10.3741-3751.1997](https://doi.org/10.1128/aem.63.10.3741-3751.1997)
 11. Schrader, C., Schielke, A., Ellerbroek, L., & Johne, R. (2012). PCR inhibitors: Occurrence, properties and removal. *Journal of Applied Microbiology*, 113(5), 1014–1026. DOI: [10.1111/j.1365-2672.2012.05384.x](https://doi.org/10.1111/j.1365-2672.2012.05384.x)
 12. Kim, C. S., Lee, C. H., Shin, J. S., Chung, Y. S., & Hyung, N. I. (1997). A simple and rapid method for isolation of high quality genomic DNA from fruit trees and conifers using PVP. *Nucleic Acids Research*, 25(5), 1085–1086. DOI: [10.1093/nar/25.5.1085](https://doi.org/10.1093/nar/25.5.1085)
 13. John, M. E. (1992). An efficient method for isolation of RNA and DNA from plants containing polyphenolics. *Nucleic Acids Research*, 20(9), 2381. DOI: [10.1093/nar/20.9.2381](https://doi.org/10.1093/nar/20.9.2381)
 14. Wilfinger, W. W., Mackey, K., & Chomczynski, P. (1997). Effect of pH and ionic strength on the spectrophotometric assessment of nucleic acid purity. *BioTechniques*, 22(3), 474–481. DOI: [10.2144/97223st01](https://doi.org/10.2144/97223st01)
 15. Al-Soud, W. A., & Rådström, P. (1998). Capacity of nine thermostable DNA polymerases to mediate DNA amplification in the presence of PCR-inhibiting samples. *Applied and Environmental Microbiology*, 64(10), 3748–3753. DOI: [10.1128/AEM.64.10.3748-3753.1998](https://doi.org/10.1128/AEM.64.10.3748-3753.1998)
 16. Paterson, A. H., Brubaker, C. L., & Wendel, J. F. (1993). A rapid method for extraction of cotton (*Gossypium*) genomic DNA suitable for RFLP or PCR analysis. *Plant Molecular Biology Reporter*, 11(2), 122–127. DOI: [10.1007/BF02670470](https://doi.org/10.1007/BF02670470)
 17. Lodhi, M. A., Ye, G.-N., Weeden, N. F., & Reisch, B. I. (1994). A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. *Plant Molecular Biology Reporter*, 12(1), 6–13. DOI: [10.1007/BF02668658](https://doi.org/10.1007/BF02668658)
 18. Dabo, S. M., Mitchell, E. D., & Melcher, U. (1993). A method for the isolation of nuclear DNA from cotton (*Gossypium*) leaves. *Analytical Biochemistry*, 210(1), 34–38. DOI: [10.1006/abio.1993.1146](https://doi.org/10.1006/abio.1993.1146)
 19. Fulton, T. M., Chunwongse, J., & Tanksley, S. D. (1995). Microprep protocol for extraction of DNA from tomato and other herbaceous plants. *Plant Molecular Biology Reporter*, 13(3), 207–209. DOI: [10.1007/BF02670897](https://doi.org/10.1007/BF02670897)
 20. Sahu, S. K., Thangaraj, M., & Kathiresan, K. (2012). DNA extraction protocol for plants with high levels of secondary metabolites and polysaccharides without using liquid nitrogen and phenol. *ISRN Molecular Biology*, 2012, 205049. DOI: [10.5402/2012/205049](https://doi.org/10.5402/2012/205049)
 21. Ahmed, I., Islam, M., Arshad, W., Mannan, A., Ahmad, W., & Mirza, B. (2009). High-quality plant DNA extraction for PCR: An easy approach. *Journal of Applied Genetics*, 50(2), 105–107. DOI: [10.1007/BF03195661](https://doi.org/10.1007/BF03195661)
 22. Allen, G. C., Flores-Vergara, M. A., Krasynanski, S., Kumar, S., & Thompson, W. F. (2006). A modified protocol for rapid DNA isolation from plant tissues using cetyltrimethylammonium bromide. *Nature Protocols*, 1(5), 2320–2325. DOI: [10.1038/nprot.2006.384](https://doi.org/10.1038/nprot.2006.384)
 23. Wang, H., Qi, M., & Cutler, A. J. (1993). A simple method of preparing plant samples for PCR. *Nucleic Acids Research*, 21(17), 4153–4154. DOI: [10.1093/nar/21.17.4153](https://doi.org/10.1093/nar/21.17.4153)
 24. Xin, Z., & Chen, J. (2012). A high throughput DNA extraction method with high yield and quality. *Plant Methods*, 8, 26. DOI: [10.1186/1746-4811-8-26](https://doi.org/10.1186/1746-4811-8-26)
 25. Chase, M. W., & Hills, H. H. (1991). Silica gel: An ideal material for field preservation of leaf samples for DNA studies. *Taxon*, 40(2), 215–220. DOI: [10.2307/1222975](https://doi.org/10.2307/1222975)



26. Rogstad, S. H. (1992). Saturated NaCl-CTAB solution as a means of field preservation of leaves for DNA analyses. *Taxon*, 41(4), 701–708. DOI: [10.2307/1222395](https://doi.org/10.2307/1222395)
27. Staats, M., Cuenca, A., Richardson, J. E., Vrieling-van Ginkel, R., Petersen, G., Seberg, O., & Bakker, F. T. (2011). DNA damage in plant herbarium tissue. *PLoS ONE*, 6(12), e28448. DOI: [10.1371/journal.pone.0028448](https://doi.org/10.1371/journal.pone.0028448)
28. Särkinen, T., Staats, M., Richardson, J. E., Cowan, R. S., & Bakker, F. T. (2012). How to open the treasure chest? Optimising DNA extraction from herbarium specimens. *PLoS ONE*, 7(8), e43808. DOI: [10.1371/journal.pone.0043808](https://doi.org/10.1371/journal.pone.0043808)
29. Koonjul, P. K., Brandt, W. F., Farrant, J. M., & Lindsey, G. G. (1999). Inclusion of polyvinylpyrrolidone in the polymerase chain reaction reverses the inhibitory effects of polyphenolic contamination of RNA. *Nucleic Acids Research*, 27(3), 915–916. DOI: [10.1093/nar/27.3.915](https://doi.org/10.1093/nar/27.3.915)
30. Desjardins, P., & Conklin, D. (2010). NanoDrop microvolume quantitation of nucleic acids. *Journal of Visualized Experiments*, 45, e2565. DOI: [10.3791/2565](https://doi.org/10.3791/2565)
31. Lee, P. Y., Costumbrado, J., Hsu, C.-Y., & Kim, Y. H. (2012). Agarose gel electrophoresis for the separation of DNA fragments. *Journal of Visualized Experiments*, 62, e3923. DOI: [10.3791/3923](https://doi.org/10.3791/3923)
32. Bustin, S. A., Benes, V., Garson, J. A., Hellemans, J., Huggett, J., Kubista, M., ... Wittwer, C. T. (2009). The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry*, 55(4), 611–622. DOI: [10.1373/clinchem.2008.112797](https://doi.org/10.1373/clinchem.2008.112797)
33. Mullis, K. B., & Faloona, F. A. (1987). Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods in Enzymology*, 155, 335–350. DOI: [10.1016/0076-6879\(87\)55023-6](https://doi.org/10.1016/0076-6879(87)55023-6)
34. Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., ... Erlich, H. A. (1988). Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*, 239(4839), 487–491. DOI: [10.1126/science.2448875](https://doi.org/10.1126/science.2448875)
35. Williams, J. G. K., Kubelik, A. R., Livak, K. J., Rafalski, J. A., & Tingey, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, 18(22), 6531–6535. DOI: [10.1093/nar/18.22.6531](https://doi.org/10.1093/nar/18.22.6531)
36. Vos, P., Hogers, R., Bleeker, M., Reijans, M., van de Lee, T., Hornes, M., ... Zabeau, M. (1995). AFLP: A new technique for DNA fingerprinting. *Nucleic Acids Research*, 23(21), 4407–4414. DOI: [10.1093/nar/23.21.4407](https://doi.org/10.1093/nar/23.21.4407)
37. CBOL Plant Working Group. (2009). A DNA barcode for land plants. *Proceedings of the National Academy of Sciences*, 106(31), 12794–12797. DOI: [10.1073/pnas.0905845106](https://doi.org/10.1073/pnas.0905845106)
38. Kress, W. J., Wurdack, K. J., Zimmer, E. A., Weigt, L. A., & Janzen, D. H. (2005). Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences*, 102(23), 8369–8374. DOI: [10.1073/pnas.0503123102](https://doi.org/10.1073/pnas.0503123102)
39. Taberlet, P., Gielly, L., Pautou, G., & Bouvet, J. (1991). Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Molecular Biology*, 17(5), 1105–1109. DOI: [10.1007/BF00037152](https://doi.org/10.1007/BF00037152)
40. Rafalski, A. (2002). Applications of single nucleotide polymorphisms in crop genetics. *Current Opinion in Plant Biology*, 5(2), 94–100. DOI: [10.1016/S1369-5266\(02\)00240-6](https://doi.org/10.1016/S1369-5266(02)00240-6)
41. Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell, S. E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE*, 6(5), e19379. DOI: [10.1371/journal.pone.0019379](https://doi.org/10.1371/journal.pone.0019379)
42. Metzker, M. L. (2010). Sequencing technologies: The next generation. *Nature Reviews Genetics*, 11(1), 31–46. DOI: [10.1038/nrg2626](https://doi.org/10.1038/nrg2626)