



ISSN : 2347 - 2243

Indo - American Journal of Life Sciences and Biotechnology



<https://iajlb.org>

Email : editor@iajlb.com or iajlb.editor@gmail.com



Genetic Diversity of Indigenous Rice Landraces Assessed Using SSR Markers

Matteo Vitale¹, Guillaume Thibaudeau², Tiago Teixeira³

¹Associate Professor, Department of Genetics, Monti Lepini Institute of Life Sciences, Frosinone, Italy.

²Senior Research Fellow, Department of Human Genetics, Vosges Institute of Applied Microbiology, Epinal, France.

³Associate Professor, Department of Human Genetics, Minho Valley Institute of Health Research, Viana do Castelo, Portugal.

Email: ¹m.vitale@mlils.it, ²g.thibaudeau@viam-e.fr, ³t.teixeira@mvihp.pt

ORCID: ¹0000-0001-1998-1062, ²0000-0001-0921-5444, ³0000-0001-2304-0644

Received: February 27, 2013 Accepted: July 25, 2013 Published: November 17, 2013

DOI: [10.62644/iajlb.2013.v10.i4.pp11-18](https://doi.org/10.62644/iajlb.2013.v10.i4.pp11-18)

ABSTRACT

Farmer-maintained rice landraces hold much of the variation that breeders and gene banks hope to use, and simple sequence repeat (SSR) markers have been the workhorse for describing it for almost two decades. This narrative review asks what an SSR survey of indigenous rice landraces can and cannot tell a reader. It draws on published work on rice domestication and population structure, on the discovery and behaviour of microsatellites, on the statistics used to summarise allelic data, and on the practical problems of sampling and scoring, and it reports no new data. SSR markers are co-dominant, multi-allelic and cheap to run, which makes them well suited to comparing many accessions at modest cost; they are also prone to scoring differences between laboratories, null alleles and size homoplasy, and a limited set of loci may say little about genes of agronomic value. Rice is largely self-pollinating, so landrace accessions are mostly homozygous and the usual heterozygosity-based summaries must be read with care. Common weaknesses in published surveys include unclear definitions of what counts as a landrace, small or uneven samples, unreported scoring rules and over-interpretation of clusters. The review concludes that SSR surveys remain a sound first step for describing and conserving landrace diversity, provided that sampling, scoring and analysis are reported openly, and it sets out a short list of reporting practices.

Keywords: *Oryza sativa*; Rice landraces; SSR markers; Microsatellites; Genetic diversity; Population structure; Narrative review

Citation: Vitale, Thibaudeau and Teixeira [2013]. Genetic Diversity of Indigenous Rice Landraces Assessed Using SSR Markers. *Indo-Am. J. Life Sci. Biotechnol.* 10(4):11-18. DOI: [10.62644/iajlb.2013.v10.i4.pp11-18](https://doi.org/10.62644/iajlb.2013.v10.i4.pp11-18)

Article Class: Narrative Review

1. Introduction

Rice feeds more people than any other crop, and much of the variation within it sits in the fields of farmers who have kept

and exchanged their own seed for generations (1, 2). These farmer-maintained varieties, usually called landraces, were shaped by local climate,



soil and cooking preference. They are also the raw material that breeders and gene banks hope to draw on when conditions change (3, 4). Describing how much variation they hold, and how it is arranged, is a practical question.

For almost two decades the usual way of asking that question has been to genotype a set of accessions with simple sequence repeat (SSR) markers, also called microsatellites, and to summarise the allelic data (5, 6, 7). The method is inexpensive, which is why many surveys of indigenous rice have used it (8, 9). It is also easy to apply carelessly.

This review sets out what is known about rice diversity and landraces, what SSR markers measure, how the data are summarised and how much weight a survey can carry. The article is a narrative review. It reports no new data and describes no collection, locality or laboratory work of the authors, and where it refers to a typical survey it means a pattern in the published literature and not any single study. The aim is to be plain about what such a survey can support.

2. Literature Review

2.1 Rice Diversity and the Landrace Concept

Asian cultivated rice is not one uniform group (Figure 1 lists some milestones in how that was learned). Isozyme work separated the varieties into several clusters long ago (10), and later work with microsatellites resolved a small number of groups that correspond broadly to indica, aus, aromatic and the two kinds of japonica (11). How those groups arose is still argued, with analyses favouring either more than one domestication or a single origin followed by gene flow (12, 13, 14, 15).

The word landrace is less settled than it looks. Reviews of the term describe a

locally adapted population with a local name, internal genetic variation and no formal breeding history (16, 17). In practice a name can cover several distinct lines, and two names can cover the same one. Surveys of farmer-maintained varieties show that the number of named varieties and their evenness in the field both matter (1). A study that does not say how its landraces were identified leaves the reader unable to judge what the diversity figure refers to. This is why one landrace can look distinct in one paper and ordinary in the next.




Figure 1. Selected milestones in the literature on rice diversity and markers (schematic timeline, not to scale).

2.2 What SSR Markers Are

A microsatellite is a short motif, commonly two to six bases, repeated in tandem. Rice carries many of them over its twelve chromosomes, and their frequency and length variation have been catalogued in detail (18, 19). Loci differ in repeat number between plants, so one PCR with flanking primers yields alleles told apart by length. Large public sets of mapped primer pairs (6, 7) explain much of why the method spread.

Compared with older marker systems, SSRs are co-dominant and multi-allelic, and they transfer easily between laboratories that share primer sequences (20, 21). Table 1 and Figure 2 set them



beside the systems that came before and after. Their strength is the information in each locus; their weakness is that mutation by slippage is fast and not always orderly (22, 23).

Table 1. Marker systems used to describe rice diversity.

Marker	Strength	Main weakness
Isozymes	Cheap; early view of rice groups (10)	Few loci; protein level
RFLP	Co-dominant; stable	Slow; needs much DNA (21)
RAPD and AFLP	Many bands without prior sequence	Dominant; repeatability (21)
SSR	Co-dominant, multi-allelic, PCR-based (19, 20)	Scoring drift; null alleles (24)
SNP	Abundant; automated calls (25)	Needs prior discovery; fewer alleles (26)

Note: Qualitative comparison drawn from the cited methodological literature; no figures are given because they depend on species, panel and laboratory.




Figure 2. Qualitative comparison of marker systems (schematic matrix, not data).

2.3 Scoring, Errors and Comparability

Scoring is where many problems begin. Stutter bands, null alleles and different sizing between runs all produce genotyping errors that bias distance and structure estimates (24). Reviews of microsatellite genotyping stress reference samples, repeated genotypes and clear

scoring rules (27, 28). Such artefacts are small in any one run, but they add up across a panel.

2.4 From Alleles to Statements About Diversity

Several summaries are standard. Allele number and richness describe variants present, and they rise with sample size unless corrected (29). Gene diversity describes the chance that two randomly chosen alleles differ (30). Differentiation between groups is expressed with a fixation index (31) or with partitioning of variance (32), and relatedness is drawn as a distance tree (33, 34). Model-based clustering infers ancestry groups from the genotypes (35), and a separate step is needed to choose how many groups to report (36). Free programs calculate most of these (37, 38, 39). Table 2 lists them with the main caution for each.

Table 2. Summaries commonly reported in SSR surveys of landraces.

Summary	What it describes	Caution
Allele number and richness	How many variants are present	Depends on sample size (29)
Gene diversity	Chance two alleles differ (30)	Heterozygote counts mislead in selfers
Fixation index	Differentiation between groups (31)	Not a true measure of differentiation (40)
Molecular variance	Split between and within groups (32)	Groups must be defined first
Distance tree	Relatedness of accessions (33, 34)	Branches may not be reliable
Model-based clusters	Inferred ancestry groups (35, 36)	Cluster number can be unstable

Note: The last column gives the authors' reading of the main caution for each summary.

Rice complicates two of these. It is mostly self-pollinating, so plants within a landrace



are largely homozygous for biological reasons, not because diversity has been lost, and heterozygosity-based quantities tell less than they would for an outcrossing crop. Differentiation measures also have mathematical limits when loci have many alleles (40). Seen this way, a low diversity value in a small sample is not evidence of a narrow gene pool.

2.5 SSRs, SNPs and Genome-Scale Data

Single nucleotide polymorphisms are now the main rival. They are abundant, can be called automatically, and have been used to relate landraces to modern varieties and to map traits (25, 41, 42). A maize comparison found that SSRs carry more information per locus, but that many SNPs can match a small SSR set for relatedness and structure (26). SSRs remain practical for laboratories without sequencing facilities and for gene bank work such as defining core subsets (43, 44).

3. Materials and Methods

3.1 Scope and Question

The review asks one question in two parts: what does an SSR survey of indigenous rice landraces measure, and how far can its results be generalised? The question was broken into the steps a survey follows (Figure 3): choosing and naming accessions, typing and scoring DNA, summarising data, and interpreting clusters. None of this is exotic, but each step is easy to skip.

3.2 Literature Assembly

The review draws on peer-reviewed work in four areas: rice domestication and population structure; the discovery and behaviour of microsatellites; the statistics used to summarise marker data; and the conservation of crop diversity. Sources were chosen for relevance and clarity of method; work on other regions and crops is used only for method.




Figure 3. Workflow of an SSR diversity survey and the point at which each common error enters (schematic, not data).

We did not run a formal systematic search and we do not claim one. This is a narrative synthesis, and the choice of sources reflects the authors' judgement. The review is therefore a better guide to how a survey should be designed and read than to the size of any diversity value.

3.3 Appraisal Framework

To judge a survey we used four short questions. Is the landrace concept defined and the source of each accession recorded (1, 16)? Is scoring described well enough to be repeated, with errors and missing data reported (24, 27)? Is the choice of summary suited to a self-pollinating crop (30, 40)? And are clusters presented with the method used to choose their number (36)? We applied the questions qualitatively, as a reading guide, and did not give any survey a score. Table 3 turns them into reporting points.

4. Results

4.1 What SSR Surveys of Landraces Typically Show

Where SSR surveys of traditional rice have been published, the broad pattern is consistent. Landrace panels usually separate into a few groups that echo the



major groups of cultivated rice (11, 45), and regional collections often hold groups that are partly distinct from those of improved varieties (8, 9). Differences in marker panels and sample design make direct comparison between surveys hard, so the shared pattern is a qualitative one.

Table 3. Reporting points for an SSR survey of landraces.

Point	Why it matters	Practice
Landrace definition	Names and types vary (16, 17)	State the working definition
Collection record	Source decides what is sampled (1)	Give place, date and source
Allele scoring	Sizes shift between runs (27)	Use reference samples and rules
Missing data and errors	Bias distance and structure (24)	Report rates and repeats
Marker choice	Locus choice shapes results (28)	List loci and positions
Cluster number	Software output is not proof (36)	Compare methods

Note: Points are drawn from the cited literature on microsatellite genotyping and on landrace definition; they are not a formal standard.

4.2 Sampling and Scoring Decisions

The conclusion of a survey depends on decisions made before the first gel or run. Sampling one plant per accession hides variation within the landrace, while sampling several raises cost and, in a selfing crop, may show near-identical lines (16). Loci spread over all twelve chromosomes give a better view than a handful from one region (7, 28).

Scoring decisions matter just as much. Rules for doubtful peaks, repeated samples and a record of missing data all reduce the chance that the result describes the instrument rather than the rice (24, 27). A survey that reports none of these has not necessarily been done badly, but the reader cannot tell.

4.3 Reading Clusters and Distances

Clusters are the most quotable result and the easiest to over-read. A model-based program returns a partition for whatever number of groups it is asked for, and the best-looking number may not make biological sense (35, 36). Trees can show tidy branches that are poorly supported, and a distance depends on the coefficient chosen (33, 34). Analysis of variance among groups needs the groups to be defined in advance (32). A cluster that disappears when the method is changed was probably never a stable feature of the material.

Put plainly, the literature supports a measured statement. SSR surveys are a sound way of sorting landraces into broad groups and of showing where diversity is concentrated; they are a weaker guide to the finer arrangement of variation, to within-landrace diversity, or to the genes that matter for adaptation. A survey that makes only the first kind of claim is on firm ground. It is a modest conclusion, and it is the one the literature will bear.

5. Discussion

5.1 Interpretation and Practical Reading

The main finding of this review is conditional. SSR data are most informative when the landrace concept is defined, the sample is balanced and the scoring is controlled, and they are least informative when none of these is true. The question for a reader is not whether SSRs work but whether a given survey did what makes them work (27, 28). Nothing here argues against SSR work; it argues for reporting what was done.

Two findings would change this reading: close agreement between SSR and SNP groupings on the same panel, and evidence that landrace names track genetic identity within a region (17).



5.2 Limitations

This is a narrative review and shares the weaknesses of that form. The selection of sources reflects judgement, no formal search or quality scoring was done, and no numerical results from individual surveys are quoted. A reader who needs a diversity value for a specific region should go to the original studies. Work on other crops, such as the maize comparison (26), is an imperfect proxy for rice landraces.

5.3 Directions for Future Work

Three needs stand out. First, shared reference samples and open scoring files. Second, surveys that sample several plants per landrace and keep the collection record (1, 16). Third, wider use of SNP panels on the same accessions, so that SSR-based groupings can be checked against genome-scale data (15, 25). It is a habit worth keeping even when the answer seems obvious.

6. Conclusion

SSR markers remain a practical and informative way to describe the diversity of indigenous rice landraces, and a first look at a collection is still sensibly made with them. What a survey can claim depends on how the landraces were defined and sampled, how the alleles were scored and how the groups were chosen.

For the practitioner the message is encouraging rather than discouraging. A careful, openly reported SSR survey can guide collection and conservation at low cost.

7. Acknowledgements

The authors thank colleagues who commented on earlier drafts of this review.

8. Funding

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

9. Conflict of Interest

The authors declare no conflict of interest.

10. Author Contributions

M.V. conceived the review and wrote the sections on rice diversity and landraces. G.T. wrote the sections on microsatellite markers and genotyping. T.T. wrote the sections on data analysis and appraisal. All authors revised the manuscript and approved the final version.

11. Ethics Statement

This is a review of published literature. No human participants or animals were used and no ethical approval was required.

12. Data Availability

No new data were generated. All sources are cited in the reference list.

13. References

1. Jarvis, D. I., Brown, A. H. D., Cuong, P. H., Collado-Panduro, L., Latournerie-Moreno, L., Gyawali, S., ... Hodgkin, T. (2008). A global perspective of the richness and evenness of traditional crop-variety diversity maintained by farming communities. *Proceedings of the National Academy of Sciences*, 105(14), 5326–5331. DOI: [10.1073/pnas.0800607105](https://doi.org/10.1073/pnas.0800607105)
2. Vaughan, D. A., Lu, B.-R., & Tomooka, N. (2008). The evolving story of rice evolution. *Plant Science*, 174(4), 394–408. DOI: [10.1016/j.plantsci.2008.01.016](https://doi.org/10.1016/j.plantsci.2008.01.016)
3. Tanksley, S. D., & McCouch, S. R. (1997). Seed banks and molecular maps: Unlocking genetic potential from the wild. *Science*, 277(5329), 1063–1066. DOI: [10.1126/science.277.5329.1063](https://doi.org/10.1126/science.277.5329.1063)
4. McCouch, S. R., McNally, K. L., Wang, W., & Sackville Hamilton, R. (2012). Genomics of gene banks: A case study in rice. *American Journal of Botany*, 99(2), 407–423. DOI: [10.3732/ajb.1100385](https://doi.org/10.3732/ajb.1100385)
5. Akagi, H., Yokozeki, Y., Inagaki, A., & Fujimura, T. (1996). Microsatellite DNA markers for rice chromosomes. *Theoretical and Applied Genetics*, 93(7), 1071–1077. DOI: [10.1007/BF00230127](https://doi.org/10.1007/BF00230127)
6. Chen, X., Temnykh, S., Xu, Y., Cho, Y. G., & McCouch, S. R. (1997). Development of a microsatellite framework map providing genome-wide coverage in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, 95(4), 553–567. DOI: [10.1007/s001220050596](https://doi.org/10.1007/s001220050596)



7. McCouch, S. R., Teytelman, L., Xu, Y., Lobos, K. B., Clare, K., Walton, M., ... Stein, L. (2002). Development and mapping of 2240 new SSR markers for rice (*Oryza sativa* L.). *DNA Research*, 9(6), 199–207. DOI: [10.1093/dnares/9.6.199](https://doi.org/10.1093/dnares/9.6.199)
8. Thomson, M. J., Septiningsih, E. M., Suwardjono, F., Santoso, T. J., Silitonga, T. S., & McCouch, S. R. (2007). Genetic diversity analysis of traditional and improved Indonesian rice (*Oryza sativa* L.) germplasm using microsatellite markers. *Theoretical and Applied Genetics*, 114(3), 559–568. DOI: [10.1007/s00122-006-0457-1](https://doi.org/10.1007/s00122-006-0457-1)
9. Das, B., Sengupta, S., Parida, S. K., Roy, B., Ghosh, M., Prasad, M., & Ghose, T. K. (2013). Genetic diversity and population structure of rice landraces from Eastern and North Eastern States of India. *BMC Genetics*, 14, 71. DOI: [10.1186/1471-2156-14-71](https://doi.org/10.1186/1471-2156-14-71)
10. Glaszmann, J. C. (1987). Isozymes and classification of Asian rice varieties. *Theoretical and Applied Genetics*, 74(1), 21–30. DOI: [10.1007/BF00290078](https://doi.org/10.1007/BF00290078)
11. Garris, A. J., Tai, T. H., Coburn, J., Kresovich, S., & McCouch, S. (2005). Genetic structure and diversity in *Oryza sativa* L. *Genetics*, 169(3), 1631–1638. DOI: [10.1534/genetics.104.035642](https://doi.org/10.1534/genetics.104.035642)
12. Londo, J. P., Chiang, Y.-C., Hung, K.-H., Chiang, T.-Y., & Schaal, B. A. (2006). Phylogeography of Asian wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated rice, *Oryza sativa*. *Proceedings of the National Academy of Sciences*, 103(25), 9578–9583. DOI: [10.1073/pnas.0603152103](https://doi.org/10.1073/pnas.0603152103)
13. Kovach, M. J., Sweeney, M. T., & McCouch, S. R. (2007). New insights into the history of rice domestication. *Trends in Genetics*, 23(11), 578–587. DOI: [10.1016/j.tig.2007.08.012](https://doi.org/10.1016/j.tig.2007.08.012)
14. Sweeney, M., & McCouch, S. (2007). The complex history of the domestication of rice. *Annals of Botany*, 100(5), 951–957. DOI: [10.1093/aob/mcm128](https://doi.org/10.1093/aob/mcm128)
15. Huang, X., Kurata, N., Wei, X., Wang, Z.-X., Wang, A., Zhao, Q., ... Han, B. (2012). A map of rice genome variation reveals the origin of cultivated rice. *Nature*, 490(7421), 497–501. DOI: [10.1038/nature11532](https://doi.org/10.1038/nature11532)
16. Zeven, A. C. (1998). Landraces: A review of definitions and classifications. *Euphytica*, 104(2), 127–139. DOI: [10.1023/A:1018683119237](https://doi.org/10.1023/A:1018683119237)
17. Camacho Villa, T. C., Maxted, N., Scholten, M., & Ford-Lloyd, B. (2005). Defining and identifying crop landraces. *Plant Genetic Resources*, 3(3), 373–384. DOI: [10.1079/PGR200591](https://doi.org/10.1079/PGR200591)
18. Temnykh, S., Park, W. D., Ayres, N., Cartinhour, S., Hauck, N., Lipovich, L., ... McCouch, S. R. (2000). Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, 100(5), 697–712. DOI: [10.1007/s001220051342](https://doi.org/10.1007/s001220051342)
19. Temnykh, S., DeClerck, G., Lukashova, A., Lipovich, L., Cartinhour, S., & McCouch, S. (2001). Computational and experimental analysis of microsatellites in rice (*Oryza sativa* L.): Frequency, length variation, transposon associations, and genetic marker potential. *Genome Research*, 11(8), 1441–1452. DOI: [10.1101/gr.184001](https://doi.org/10.1101/gr.184001)
20. Morgante, M., & Olivieri, A. M. (1993). PCR-amplified microsatellites as markers in plant genetics. *The Plant Journal*, 3(1), 175–182. DOI: [10.1111/j.1365-3113X.1993.tb00020.x](https://doi.org/10.1111/j.1365-3113X.1993.tb00020.x)
21. Powell, W., Morgante, M., Andre, C., Hanafey, M., Vogel, J., Tingey, S., & Rafalski, A. (1996). The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Molecular Breeding*, 2(3), 225–238. DOI: [10.1007/BF00564200](https://doi.org/10.1007/BF00564200)
22. Ellegren, H. (2004). Microsatellites: Simple sequences with complex evolution. *Nature Reviews Genetics*, 5(6), 435–445. DOI: [10.1038/nrg1348](https://doi.org/10.1038/nrg1348)
23. Schlötterer, C. (2004). The evolution of molecular markers — just a matter of fashion? *Nature Reviews Genetics*, 5(1), 63–73. DOI: [10.1038/nrg1249](https://doi.org/10.1038/nrg1249)
24. Pompanon, F., Bonin, A., Bellemain, E., & Taberlet, P. (2005). Genotyping errors: Causes, consequences and solutions. *Nature Reviews Genetics*, 6(11), 847–859. DOI: [10.1038/nrg1707](https://doi.org/10.1038/nrg1707)
25. McNally, K. L., Childs, K. L., Bohnert, R., Davidson, R. M., Zhao, K., Ulat, V. J., ... Leach, J. E. (2009). Genomewide SNP variation reveals relationships among landraces and modern varieties of rice. *Proceedings of the National Academy of Sciences*, 106(30), 12273–12278. DOI: [10.1073/pnas.0900992106](https://doi.org/10.1073/pnas.0900992106)
26. Hamblin, M. T., Warburton, M. L., & Buckler, E. S. (2007). Empirical comparison of simple sequence repeats and single nucleotide polymorphisms in assessment of maize diversity and relatedness. *PLoS ONE*, 2(12), e1367. DOI: [10.1371/journal.pone.0001367](https://doi.org/10.1371/journal.pone.0001367)
27. Guichoux, E., Lagache, L., Wagner, S., Chaumeil, P., Léger, P., Lepais, O., ... Petit, R. J. (2011). Current trends in microsatellite genotyping. *Molecular Ecology Resources*, 11(4), 591–611. DOI: [10.1111/j.1755-0998.2011.03014.x](https://doi.org/10.1111/j.1755-0998.2011.03014.x)
28. Selkoe, K. A., & Toonen, R. J. (2006). Microsatellites for ecologists: A practical guide to using and evaluating microsatellite markers. *Ecology Letters*, 9(5), 615–629. DOI: [10.1111/j.1461-0248.2006.00889.x](https://doi.org/10.1111/j.1461-0248.2006.00889.x)



29. Kalinowski, S. T. (2005). HP-RARE 1.0: A computer program for performing rarefaction on measures of allelic richness. *Molecular Ecology Notes*, 5(1), 187–189. DOI: [10.1111/j.1471-8286.2004.00845.x](https://doi.org/10.1111/j.1471-8286.2004.00845.x)
30. Nei, M. (1973). Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences*, 70(12), 3321–3323. DOI: [10.1073/pnas.70.12.3321](https://doi.org/10.1073/pnas.70.12.3321)
31. Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population structure. *Evolution*, 38(6), 1358–1370. DOI: [10.2307/2408641](https://doi.org/10.2307/2408641)
32. Excoffier, L., Smouse, P. E., & Quattro, J. M. (1992). Analysis of molecular variance inferred from metric distances among DNA haplotypes: Application to human mitochondrial DNA restriction data. *Genetics*, 131(2), 479–491. DOI: [10.1093/genetics/131.2.479](https://doi.org/10.1093/genetics/131.2.479)
33. Nei, M., & Li, W. H. (1979). Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences*, 76(10), 5269–5273. DOI: [10.1073/pnas.76.10.5269](https://doi.org/10.1073/pnas.76.10.5269)
34. Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406–425. DOI: [10.1093/oxfordjournals.molbev.a040454](https://doi.org/10.1093/oxfordjournals.molbev.a040454)
35. Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155(2), 945–959. DOI: [10.1093/genetics/155.2.945](https://doi.org/10.1093/genetics/155.2.945)
36. Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology*, 14(8), 2611–2620. DOI: [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x)
37. Peakall, R., & Smouse, P. E. (2006). GENALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, 6(1), 288–295. DOI: [10.1111/j.1471-8286.2005.01155.x](https://doi.org/10.1111/j.1471-8286.2005.01155.x)
38. Liu, K., & Muse, S. V. (2005). PowerMarker: An integrated analysis environment for genetic marker analysis. *Bioinformatics*, 21(9), 2128–2129. DOI: [10.1093/bioinformatics/bti282](https://doi.org/10.1093/bioinformatics/bti282)
39. Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28(10), 2731–2739. DOI: [10.1093/molbev/msr121](https://doi.org/10.1093/molbev/msr121)
40. Jost, L. (2008). G_{ST} and its relatives do not measure differentiation. *Molecular Ecology*, 17(18), 4015–4026. DOI: [10.1111/j.1365-294X.2008.03887.x](https://doi.org/10.1111/j.1365-294X.2008.03887.x)
41. Huang, X., Wei, X., Sang, T., Zhao, Q., Feng, Q., Zhao, Y., ... Han, B. (2010). Genome-wide association studies of 14 agronomic traits in rice landraces. *Nature Genetics*, 42(11), 961–967. DOI: [10.1038/ng.695](https://doi.org/10.1038/ng.695)
42. Zhao, K., Tung, C.-W., Eizenga, G. C., Wright, M. H., Ali, M. L., Price, A. H., ... McCouch, S. R. (2011). Genome-wide association mapping reveals a rich genetic architecture of complex traits in *Oryza sativa*. *Nature Communications*, 2, 467. DOI: [10.1038/ncomms1467](https://doi.org/10.1038/ncomms1467)
43. Brown, A. H. D. (1989). Core collections: A practical approach to genetic resources management. *Genome*, 31(2), 818–824. DOI: [10.1139/g89-144](https://doi.org/10.1139/g89-144)
44. Ebana, K., Kojima, Y., Fukuoka, S., Nagamine, T., & Kawase, M. (2008). Development of mini core collection of Japanese rice landrace. *Breeding Science*, 58(3), 281–291. DOI: [10.1270/jsbbs.58.281](https://doi.org/10.1270/jsbbs.58.281)
45. Jain, S., Jain, R. K., & McCouch, S. R. (2004). Genetic analysis of Indian aromatic and quality rice (*Oryza sativa* L.) germplasm using panels of fluorescent-labeled microsatellite markers. *Theoretical and Applied Genetics*, 109(5), 965–977. DOI: [10.1007/s00122-004-1700-6](https://doi.org/10.1007/s00122-004-1700-6)