



ISSN : 2347 - 2243

Indo - American Journal of Life Sciences and Biotechnology



<https://iajlb.org>

Email : editor@iajlb.com or iajlb.editor@gmail.com



Expression Analysis of Drought-Responsive Genes in Contrasting Rice Genotypes Using Quantitative Real-Time PCR

Lucie Vesely¹●, Romain Rousselot²●, Ondrej Novotny³●

¹Senior Scientist, Department of Biotechnology, Bohemian Forest Institute of Bioinformatics, Klatovy, Czech Republic.

²Research Associate, Department of Biotechnology, Vosges Institute of Applied Microbiology, Epinal, France.

³Senior Research Fellow, Department of Molecular Biology, Bohemian Forest Institute of Bioinformatics, Klatovy, Czech Republic.

Email: ¹l.vesely@bfib.cz, ²r.rousselot@viam-e.fr, ³o.novotny@bfib.cz

ORCID: ¹0000-0001-3231-5419, ²0000-0001-4926-7847, ³0000-0001-4195-3535

Received: July 8, 2013 Accepted: October 26, 2013 Published: February 23, 2014

DOI: [10.62644/iajlb.2014.v11.i1.pp1-10](https://doi.org/10.62644/iajlb.2014.v11.i1.pp1-10)

ABSTRACT

Rice loses yield quickly when water runs short, and the genes that respond to drought are often studied by quantitative real-time PCR (qPCR) in genotypes chosen because they behave differently in the field. This narrative review asks what such an expression analysis can and cannot tell a reader, and what separates a convincing comparison of contrasting rice genotypes from a fragile one. It draws on published work on the signalling and regulatory networks that respond to water deficit, on transcript profiling in tolerant and sensitive rice, and on the methodology of qPCR, and it reports no new expression data. Regulators of the DREB, bZIP and NAC families, protective proteins and osmolyte pathways recur across the literature, but the direction and size of a change depend on tissue, developmental stage, severity and speed of stress, and the genotype pair. The review argues that normalisation to a reference gene that was never tested under the stress in question is the commonest and least visible source of error, followed by uneven RNA quality, unexamined amplification efficiency and treatment of technical wells as independent samples. Differential expression between genotypes is also an association, not proof that a gene causes tolerance. The review closes with a short checklist of design, reporting and interpretation points for studies of this type.

Keywords: Rice; Drought stress; Gene expression; Quantitative real-time PCR; Reference genes; Genotypes; Narrative review

Citation: Vesely, Rousselot and Novotny [2014]. Expression Analysis of Drought-Responsive Genes in Contrasting Rice Genotypes Using Quantitative Real-Time PCR. *Indo-Am. J. Life Sci. Biotechnol.* 11(1):1-10. DOI: [10.62644/iajlb.2014.v11.i1.pp1-10](https://doi.org/10.62644/iajlb.2014.v11.i1.pp1-10)

Article Class: Narrative Review

1. Introduction

Rice feeds a very large part of humanity, and it is also one of the most water-hungry cereals. Much of it is grown flooded, which makes a shortage of rain or

irrigation water unusually costly, and breeders have long looked for genotypes that hold their yield when water becomes scarce (1, 2). Some of the differences between tolerant and sensitive lines are



anatomical or physiological, for example in rooting and in the way stomata respond. Others lie in how genes are switched on and off, and that is where expression analysis enters.

Quantitative real-time PCR, usually shortened to qPCR, is the workhorse for measuring the abundance of a chosen transcript in a chosen sample (3, 4). A typical study grows two or more contrasting rice genotypes, withholds water, collects leaves or roots at intervals, and reports how a panel of candidate genes changes relative to the well-watered control. The method is sensitive and fast. It is also unforgiving of small errors in handling, because the quantity reported is a ratio of two noisy measurements. The aim here is a fair account of what the method can show and what it only appears to show.

The title of this article describes exactly that kind of experiment. This review does not report one. No plants were grown, no RNA was extracted and no expression values are presented here; the aim is to describe, from published work, what drought-responsive gene expression in rice is known to look like, how qPCR is meant to be applied to it, and which design choices decide whether a comparison between genotypes can be trusted. Readers looking for numbers from a particular study will not find them in these pages.

2. Literature Review

2.1 Drought Signalling and Gene Networks in Plants

Plants sense water deficit through changes in cell turgor, in membrane properties and in the hormone abscisic acid, and these cues are converted into changes in gene expression through several partly overlapping routes (5). Some routes depend on abscisic acid and some do not. The picture that emerged around the turn

of the century, and which later reviews refined, is of regulatory networks in which a small number of transcription factors control large sets of downstream genes (6, 7). Expression studies in rice generally try to read these networks at one or two points.

The downstream genes are a mixed group. Some encode protective proteins and enzymes that detoxify reactive oxygen species, some encode enzymes of sugar and osmolyte metabolism, and others encode transporters and water channels. Older but still useful accounts organise the response around adaptation at the level of the cell, arguing that survival under stress and productivity under stress are not the same trait (8, 9). A plant that has been stressed slowly behaves differently from one that has been stressed abruptly.

2.2 Regulators and Effectors Studied in Rice

In rice, members of the DREB family were among the first regulators shown to respond to drought, high salt and cold, with different members preferring different stresses (10). Basic leucine zipper factors such as OsbZIP23 link abscisic acid sensitivity to drought and salinity tolerance, and overexpressing it improved tolerance in transgenic plants (11). A NAC-type factor, when overexpressed, increased drought resistance and salt tolerance in rice (12). Table 1 lists the main groups and the kind of role each is expected to play.

Effector genes have been tested in a more applied way. A late embryogenesis abundant protein gene from barley, HVA1, conferred tolerance to water deficit and salt stress when expressed in transgenic rice (13), and plants engineered to accumulate trehalose showed higher tolerance to several abiotic stresses (14). General reviews of rice stress biology bring these examples together and stress



that tolerance is built from many small contributions rather than from a single switch (16). Such differences are small one at a time, but they accumulate across a whole panel of genes.

Table 1. Groups of drought-responsive genes in rice that expression studies commonly target.

Group	Examples	Usual role
Transcription factors	DREB, bZIP and NAC members (10, 11, 12)	Switch on sets of stress genes
Protective proteins	LEA proteins such as HVA1 (13)	Shield membranes and enzymes
Osmolyte pathways	Trehalose and sugar metabolism (14)	Adjust cell water balance
Signalling components	Kinases and ABA-linked factors (5, 7)	Relay the stress signal
Photosynthesis and growth genes	Changes often reported (15)	Growth traded against survival

Note: Qualitative summary. Membership of a group does not predict the direction of change in a given tissue or genotype.

2.3 Transcript Profiling in Contrasting Rice Genotypes

Microarray and related profiling studies made it possible to ask which genes respond to drought across the whole genome. Early work on cold, drought, high salinity and abscisic acid identified overlapping sets of rice genes that responded to more than one treatment (17), and work on the initial phase of salt stress showed how quickly the transcript profile of rice can shift (18). For drought itself, several groups compared cultivars that differ in tolerance. One study profiled rice cultivars exposed to long-term drought (15), another compared indica genotypes with contrasting tolerance (19), and a third followed temporal and spatial patterns across tissues (20).

These studies agree on the broad outline: many genes change, a core set is shared between genotypes, and tolerant lines often differ from sensitive ones in timing or in the strength of the response rather than in an on-off pattern. They disagree, and this matters, about particular genes. A transcript that rises in one tissue may fall in another, and a gene that separates two genotypes at one stage may not separate them at the next. This is why the same gene can look decisive in one report and unremarkable in the next.

2.4 Principles of Quantitative Real-Time PCR

The principle is simple. Fluorescence from a reporter rises as a target is copied, and the cycle at which it crosses a threshold falls earlier when more starting template is present. In practice the measurement is a chain: RNA extraction, reverse transcription to complementary DNA, amplification, detection and calculation (21, 22). Each link can bias the result. Extraction methods based on guanidinium and phenol remain standard (23), but the integrity of the RNA that comes out affects the performance of the assay and should be checked rather than assumed (24).


Relative quantification then compares a target with a reference. The widely used comparative threshold approach assumes that the target and the reference amplify with the same, near-ideal efficiency (25, 26). When efficiencies differ, a model that corrects for them gives a more honest ratio (27), and efficiency can be estimated from the amplification curves themselves rather than from a separate dilution series (28, 29). Figure 1 sketches the full chain and the point at which each safeguard belongs.

2.5 Normalisation and Reference Genes

Normalisation corrects for differences in the amount and quality of material put into each reaction. The usual device is a



reference gene thought to be expressed at a constant level. The difficulty is that no gene is constant under every condition, and a gene that was stable across tissues may not be stable under stress (30, 31). Plant studies make the point repeatedly: stability has to be shown for the species, the tissue and the treatment actually used (32, 33).



Schematic, not data

Figure 1. From stressed plant to interpretable expression value, with the checks that surround each step (schematic, not data).

For rice, validation work found that several commonly used housekeeping genes were less steady than others, and that the best choice depended on tissue and stress (34, 35). Later work mined transcriptome datasets for rice genes that are stable across organs, developmental stages and stress treatments (36), an approach that had already proved useful in *Arabidopsis* (37) and that has parallels in potato under stress (38). Software that ranks candidates, including geNorm, NormFinder, BestKeeper and qBase, helps choose among them (39, 40, 41, 42). Table 2 sets these strategies side by side. A short pilot run on the actual samples is cheap insurance against drift once leaves lose water.

Table 2. Normalisation strategies used with qPCR and their main weakness.

Strategy	What it offers	Main weakness
Single classical reference gene	Simple and cheap (35)	May itself change under stress (34)
Several validated genes	Averaged signal is steadier (39)	Needs a pilot experiment
Model-based ranking	Weights variation within and between groups (40)	Assumes the set is not co-regulated
Pair-wise correlation tools	Quick screen of candidates (41)	Sensitive to outlying samples
Reference genes from transcriptome data	Candidates chosen on stability (36, 37)	Still needs checking in your own samples (32)

Note: Strengths and weaknesses are those discussed in the cited methodological papers.

3. Materials and Methods

3.1 Scope and Question

This is a narrative review. Its question is narrow: how should an expression analysis of drought-responsive genes in contrasting rice genotypes by qPCR be designed, read and reported so that its conclusions are not larger than its evidence? The review is concerned with the logic of such studies and not with any particular dataset, and it contains no experimental work of its own.

3.2 Literature Assembly

The sources are papers the authors already knew or located through ordinary reading and citation tracing. There was no formal systematic search, no protocol registered in advance and no screening in duplicate, so the review may have missed relevant work, and the selection reflects the judgement of the authors. Only publications that appeared in or before 2014 are cited. Preference was given to primary studies of rice, to widely used methodological papers



and to reviews that organise the field. Reviews of this kind are more useful when they say where the evidence stops than when they smooth it over.

3.3 Appraisal Framework


Each source was read for four things: how the stress was imposed and monitored, how genotypes were chosen and compared, how expression was normalised, and how far the stated conclusion went beyond the measurements. Methodological sources were read against the published reporting guidance for qPCR experiments (43, 44). No scoring system or numerical weighting was applied. The appraisal is qualitative and is presented in the tables and figures that follow.

4. Results

4.1 What Drought-Responsive Expression Looks Like in Rice

Across the sources, a handful of patterns recur. Regulatory genes of the DREB, bZIP and NAC types tend to respond early, while genes for protective proteins and metabolic enzymes follow, which gives the layered picture shown in Figure 2 (6, 7). Genes tied to photosynthesis and growth are frequently reported to change as the plant shifts from growth towards maintenance (9, 15). The direction of a change is not fixed, and a given gene can rise in roots while falling in leaves (20).

Tolerance in a genotype pair is rarely explained by a single gene. Comparisons of contrasting indica lines point to differences in the number of responsive genes, in the timing of induction, and in baseline levels before stress begins (15, 19). A higher baseline can be misread as a stronger response if only the stressed samples are compared, which is one reason well-watered controls for each genotype are indispensable.



Schematic, not data

Figure 2. Layered view of a drought response, from signal to phenotype, and what an expression study samples (schematic, not data).

4.2 How the Genotype Contrast Is Built

The word contrasting hides a good deal. Two genotypes may differ in rooting depth, in stomatal behaviour, in leaf rolling or in flowering time, and the physiological basis of their tolerance may differ from the genes being measured (1, 2). Genotypes also differ in how quickly the soil dries around them and how fast the leaves lose water, so the same number of days without watering does not impose the same stress. Sampling at matched leaf water status rather than matched days removes one source of confusion (9). A reader of an expression paper is entitled to ask a plain question: would this result survive a repeat with new plants?

Related to this, the expression difference between a tolerant and a sensitive line is an association. It may reflect a cause of tolerance, a consequence of it, or simply the fact that the sensitive plant is more stressed at the moment of sampling. Transgenic studies, in which a regulator or effector gene is introduced and the plant becomes more tolerant, give stronger evidence of causation (12, 13, 14), though even those are mostly judged under



controlled conditions and not in farmers' fields.

4.3 Technical Sources of Error

The technical literature identifies a short list of recurring problems, collected in [Table 3](#). RNA of poor or uneven quality changes apparent expression in a way that is not corrected by dividing by a reference gene ([24](#)). Reverse transcription is itself variable, and including controls without the enzyme shows whether genomic DNA is contributing a signal ([21](#)). Assays for different genes amplify with different efficiency, so a single assumed value is rarely safe ([27, 29](#)). None of these steps is difficult, but each one is easy to skip when a deadline is near.

Table 3. Where a genotype comparison by qPCR can go wrong, and how to guard against it.

Step	Risk	Safeguard
Sampling	Genotypes differ in water status at the same time point (9)	Match by stress level, not clock time
RNA extraction	Degraded or impure RNA (23, 24)	Check integrity; same method for all
Reverse transcription	Uneven priming and yield (21)	Include no-enzyme controls
Amplification	Efficiency differs between assays (27, 29)	Measure efficiency per assay
Normalisation	Unstable reference gene (31)	Validate under the stress used
Statistics	Technical wells counted as replicates	Biological replicates as the unit

Note: Compiled from the cited methodological literature; it is not a reanalysis of any study.

Replication is the other weak point. Technical replicates, meaning repeated wells from the same cDNA, reduce pipetting noise but say nothing about variation between plants. The unit that

supports a claim about a genotype is the independent biological sample, and each such sample should be handled separately from stress treatment onwards ([33, 44](#)). Treating wells as if they were plants gives confidence intervals that are far too narrow.

4.4 The Normalisation Problem in Practice

Perhaps the most consequential finding of the methodological literature is how often normalisation goes unexamined. Reviews of plant qPCR papers have repeatedly noted that reference genes are chosen by habit and not validated for the conditions used ([31, 32](#)). In a drought experiment this is a serious matter, because water deficit can alter the expression of genes that were taken to be constant, and the effect on the reported fold change can be large enough to reverse its apparent direction ([34](#)).

Using more than one reference gene and checking their agreement across all treatment and genotype groups is the usual remedy ([30, 39](#)). If the chosen genes disagree, the disagreement is itself informative. [Figure 3](#) shows, in conceptual form, how a transcript difference should be read according to whether the normalisation has been checked and whether the genotype pair was matched for stress. A fold change that vanishes when the reference gene is swapped is not a finding a colleague could build on.

5. Discussion

5.1 Interpretation and Practical Reading

Read as a whole, the literature supports a measured position. Drought-responsive expression in rice is real, reproducible in outline, and informative about the biology of tolerance. But the specific numbers in any single qPCR comparison between two genotypes depend heavily on choices that are often invisible in the abstract: the



reference gene, the stress protocol, the tissue and the time point. That is a modest claim, and it is the one the literature will currently bear. The point is not that a study is poor, only that its headline travels less far than its abstract suggests.

	Reference genes validated	Reference genes not validated
Stress matched	Difference reasonably firm, still an association	Size and direction uncertain
Stress not matched	May reflect unequal stress level	Treat as a lead only

Figure 3. How the reading of a transcript difference changes with the normalisation check and the genotype pair (conceptual matrix, not data).

For a reader, the practical consequence is to ask a few questions of any paper in this area. Was the stress measured, or only imposed? Were reference genes validated under that stress? Were biological replicates independent? Were amplification efficiencies reported? Do the conclusions refer to genes measured, or do they slip into claims about tolerance in the field? Table 3 can serve as a checklist for these questions. It is a habit worth keeping even when the result looks exactly as expected.

5.2 Limitations

The review has limits that readers should keep in mind. It is narrative and not systematic, and its selection of papers cannot be called exhaustive. It is restricted to work published up to 2014, so later developments in high-throughput sequencing of transcripts and in new analysis tools are not covered. Many of the studies cited examine single cultivars or small sets, and their findings may not

extend to other genetic backgrounds or growing conditions. Because drought experiments differ in severity, duration and soil, results from pots, hydroponic systems and fields are not interchangeable. Read this way, the absence of a difference in a small experiment is not evidence that no difference exists.

There is also a limit built into the topic. A review of methods cannot say what the expression pattern of any particular genotype pair will be. Statements here about direction and timing of expression are general tendencies drawn from published work and should not be applied to a specific line without measurement.

5.3 Directions for Future Work

Three needs stand out. First, reference-gene validation should be treated as part of the experiment and reported as such, ideally with the data that led to the choice. Second, genotype comparisons would be more informative if stress were matched on physiological status and if sampling covered several time points, because a single time point confounds the size of the response with its timing. Third, expression findings would carry more weight if paired with measurements at the protein or phenotype level, since transcript abundance is not the same as function.

Finally, wider adoption of reporting guidance would help readers judge studies and repeat them (43, 44). Full reporting costs a few extra lines in a methods section and spares others the work of guessing. These points apply with equal force to any stress and any crop, not only to rice under drought.

6. Conclusion

Expression analysis of drought-responsive genes by qPCR is a powerful way to study how contrasting rice genotypes cope with water deficit, provided its limitations are respected. The regulators and effectors



most often examined are well described, but their behaviour depends on tissue, timing and the genotype pair, and so a single comparison seldom settles a question. The most common threats to a sound conclusion are an untested reference gene, uneven RNA quality, unexamined amplification efficiency, technical wells treated as replicates and the reading of association as cause. A study that addresses these openly, and that reports its methods fully, gives other researchers something they can use. Nothing in this reading argues against qPCR; it argues for using it with the care it was designed to receive.

7. Acknowledgements

The authors thank colleagues who commented on earlier versions of this review.

8. Funding

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

9. Conflict of Interest

The authors declare no conflict of interest.

10. Author Contributions

L.V. conceived the review and wrote the sections on drought signalling and rice genotypes. R.R. wrote the sections on qPCR methodology and normalisation. O.N. wrote the sections on appraisal and interpretation. All authors revised the manuscript and approved the final version.

11. Ethics Statement

This is a review of published literature. No human participants or animals were involved, and no ethical approval was required. No field trials or patient data were involved, and only published sources were used.

12. Data Availability

No new data were generated. All sources are cited in the reference list.

13. References

- Bernier, J., Atlin, G. N., Serraj, R., Kumar, A., & Spaner, D. (2008). Breeding upland rice for drought resistance. *Journal of the Science of Food and Agriculture*, 88(6), 927–939. DOI: [10.1002/jsfa.3153](https://doi.org/10.1002/jsfa.3153)
- Price, A. H., Cairns, J. E., Horton, P., Jones, H. G., & Griffiths, H. (2002). Linking drought-resistance mechanisms to drought avoidance in upland rice using a QTL approach: Progress and new opportunities to integrate stomatal and mesophyll responses. *Journal of Experimental Botany*, 53(371), 989–1004. DOI: [10.1093/jexbot/53.371.989](https://doi.org/10.1093/jexbot/53.371.989)
- Bustin, S. A. (2000). Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *Journal of Molecular Endocrinology*, 25(2), 169–193. DOI: [10.1677/jme.0.0250169](https://doi.org/10.1677/jme.0.0250169)
- Wong, M. L., & Medrano, J. F. (2005). Real-time PCR for mRNA quantitation. *BioTechniques*, 39(1), 75–85. DOI: [10.2144/05391RV01](https://doi.org/10.2144/05391RV01)
- Zhu, J.-K. (2002). Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology*, 53, 247–273. DOI: [10.1146/annurev.arplant.53.091401.143329](https://doi.org/10.1146/annurev.arplant.53.091401.143329)
- Yamaguchi-Shinozaki, K., & Shinozaki, K. (2006). Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses. *Annual Review of Plant Biology*, 57, 781–803. DOI: [10.1146/annurev.arplant.57.032905.105444](https://doi.org/10.1146/annurev.arplant.57.032905.105444)
- Shinozaki, K., & Yamaguchi-Shinozaki, K. (2007). Gene networks involved in drought stress response and tolerance. *Journal of Experimental Botany*, 58(2), 221–227. DOI: [10.1093/jxb/erl164](https://doi.org/10.1093/jxb/erl164)
- Bohnert, H. J., Nelson, D. E., & Jensen, R. G. (1995). Adaptations to environmental stresses. *The Plant Cell*, 7(7), 1099–1111. DOI: [10.1105/tpc.7.7.1099](https://doi.org/10.1105/tpc.7.7.1099)
- Farooq, M., Wahid, A., Kobayashi, N., Fujita, D., & Basra, S. M. A. (2009). Plant drought stress: Effects, mechanisms and management. *Agronomy for Sustainable Development*, 29(1), 185–212. DOI: [10.1051/agro:2008021](https://doi.org/10.1051/agro:2008021)
- Dubouzet, J. G., Sakuma, Y., Ito, Y., Kasuga, M., Dubouzet, E. G., Miura, S., ... Yamaguchi-Shinozaki, K. (2003). *OsDREB* genes in rice, *Oryza sativa* L., encode transcription activators that function in drought-, high-salt- and cold-responsive gene expression. *The Plant Journal*, 33(4), 751–763. DOI: [10.1046/j.1365-3113.2003.02111.x](https://doi.org/10.1046/j.1365-3113.2003.02111.x)



- [10.1046/j.1365-313X.2003.01661.x](https://doi.org/10.1046/j.1365-313X.2003.01661.x)
11. Xiang, Y., Tang, N., Du, H., Ye, H., & Xiong, L. (2008). Characterization of OsbZIP23 as a key player of the basic leucine zipper transcription factor family for conferring abscisic acid sensitivity and salinity and drought tolerance in rice. *Plant Physiology*, 148(4), 1938–1952. DOI: [10.1104/pp.108.128199](https://doi.org/10.1104/pp.108.128199)
 12. Hu, H., Dai, M., Yao, J., Xiao, B., Li, X., Zhang, Q., & Xiong, L. (2006). Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice. *Proceedings of the National Academy of Sciences*, 103(35), 12987–12992. DOI: [10.1073/pnas.0604882103](https://doi.org/10.1073/pnas.0604882103)
 13. Xu, D., Duan, X., Wang, B., Hong, B., Ho, T.-H. D., & Wu, R. (1996). Expression of a late embryogenesis abundant protein gene, *HVA1*, from barley confers tolerance to water deficit and salt stress in transgenic rice. *Plant Physiology*, 110(1), 249–257. DOI: [10.1104/pp.110.1.249](https://doi.org/10.1104/pp.110.1.249)
 14. Garg, A. K., Kim, J.-K., Owens, T. G., Ranwala, A. P., Choi, Y. D., Kochian, L. V., & Wu, R. J. (2002). Trehalose accumulation in rice plants confers high tolerance levels to different abiotic stresses. *Proceedings of the National Academy of Sciences*, 99(25), 15898–15903. DOI: [10.1073/pnas.252637799](https://doi.org/10.1073/pnas.252637799)
 15. Degenkolbe, T., Do, P. T., Zuther, E., Reipsilber, D., Walther, D., Hinch, D. K., & Köhl, K. I. (2009). Expression profiling of rice cultivars differing in their tolerance to long-term drought stress. *Plant Molecular Biology*, 69(1–2), 133–153. DOI: [10.1007/s11103-008-9412-7](https://doi.org/10.1007/s11103-008-9412-7)
 16. Gao, J.-P., Chao, D.-Y., & Lin, H.-X. (2007). Understanding abiotic stress tolerance mechanisms: Recent studies on stress response in rice. *Journal of Integrative Plant Biology*, 49(6), 742–750. DOI: [10.1111/j.1744-7909.2007.00495.x](https://doi.org/10.1111/j.1744-7909.2007.00495.x)
 17. Rabbani, M. A., Maruyama, K., Abe, H., Khan, M. A., Katsura, K., Ito, Y., ... Shinozaki, K. (2003). Monitoring expression profiles of rice genes under cold, drought, and high-salinity stresses and abscisic acid application using cDNA microarray and RNA gel-blot analyses. *Plant Physiology*, 133(4), 1755–1767. DOI: [10.1104/pp.103.025742](https://doi.org/10.1104/pp.103.025742)
 18. Kawasaki, S., Borchert, C., Deyholos, M., Wang, H., Brazille, S., Kawai, K., ... Bohnert, H. J. (2001). Gene expression profiles during the initial phase of salt stress in rice. *The Plant Cell*, 13(4), 889–905. DOI: [10.1105/tpc.13.4.889](https://doi.org/10.1105/tpc.13.4.889)
 19. Lenka, S. K., Katiyar, A., Chinnusamy, V., & Bansal, K. C. (2011). Comparative analysis of drought-responsive transcriptome in Indica rice genotypes with contrasting drought tolerance. *Plant Biotechnology Journal*, 9(3), 315–327. DOI: [10.1111/j.1467-7652.2010.00560.x](https://doi.org/10.1111/j.1467-7652.2010.00560.x)
 20. Wang, D., Pan, Y., Zhao, X., Zhu, L., Fu, B., & Li, Z. (2011). Genome-wide temporal-spatial gene expression profiling of drought responsiveness in rice. *BMC Genomics*, 12, 149. DOI: [10.1186/1471-2164-12-149](https://doi.org/10.1186/1471-2164-12-149)
 21. Nolan, T., Hands, R. E., & Bustin, S. A. (2006). Quantification of mRNA using real-time RT-PCR. *Nature Protocols*, 1(3), 1559–1582. DOI: [10.1038/nprot.2006.236](https://doi.org/10.1038/nprot.2006.236)
 22. Gachon, C., Mingam, A., & Charrier, B. (2004). Real-time PCR: What relevance to plant studies? *Journal of Experimental Botany*, 55(402), 1445–1454. DOI: [10.1093/jxb/erh181](https://doi.org/10.1093/jxb/erh181)
 23. Chomczynski, P., & Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry*, 162(1), 156–159. DOI: [10.1016/0003-2697\(87\)90021-2](https://doi.org/10.1016/0003-2697(87)90021-2)
 24. Fleige, S., & Pfaffl, M. W. (2006). RNA integrity and the effect on the real-time qRT-PCR performance. *Molecular Aspects of Medicine*, 27(2–3), 126–139. DOI: [10.1016/j.mam.2005.12.003](https://doi.org/10.1016/j.mam.2005.12.003)
 25. Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, 25(4), 402–408. DOI: [10.1006/meth.2001.1262](https://doi.org/10.1006/meth.2001.1262)
 26. Schmittgen, T. D., & Livak, K. J. (2008). Analyzing real-time PCR data by the comparative C_t method. *Nature Protocols*, 3(6), 1101–1108. DOI: [10.1038/nprot.2008.73](https://doi.org/10.1038/nprot.2008.73)
 27. Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, 29(9), e45. DOI: [10.1093/nar/29.9.e45](https://doi.org/10.1093/nar/29.9.e45)
 28. Ramakers, C., Ruijter, J. M., Deprez, R. H. L., & Moorman, A. F. M. (2003). Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neuroscience Letters*, 339(1), 62–66. DOI: [10.1016/S0304-3940\(02\)01423-4](https://doi.org/10.1016/S0304-3940(02)01423-4)
 29. Ruijter, J. M., Ramakers, C., Hoogaars, W. M. H., Karlen, Y., Bakker, O., van den Hoff, M. J. B., & Moorman, A. F. M. (2009). Amplification efficiency: Linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Research*, 37(6), e45. DOI: [10.1093/nar/gkp045](https://doi.org/10.1093/nar/gkp045)
 30. Huggett, J., Dheda, K., Bustin, S., & Zumla, A. (2005). Real-time RT-PCR normalisation; strategies and considerations. *Genes and Immunity*, 6(4), 279–284. DOI: [10.1038/sj.gene.6364190](https://doi.org/10.1038/sj.gene.6364190)
 31. Gutierrez, L., Mauriat, M., Guénin, S., Pelloux, J., Lefebvre, J.-F., Louvet, R., ... Bellini, C. (2008). The lack of a systematic validation of reference genes: A serious pitfall undervalued in reverse transcription-polymerase chain reaction



- (RT-PCR) analysis in plants. *Plant Biotechnology Journal*, 6(6), 609–618. DOI: [10.1111/j.1467-7652.2008.00346.x](https://doi.org/10.1111/j.1467-7652.2008.00346.x)
32. Guénin, S., Mauriat, M., Pelloux, J., Van Wuytswinkel, O., Bellini, C., & Gutierrez, L. (2009). Normalization of qRT-PCR data: The necessity of adopting a systematic, experimental conditions-specific, validation of references. *Journal of Experimental Botany*, 60(2), 487–493. DOI: [10.1093/jxb/ern305](https://doi.org/10.1093/jxb/ern305)
 33. Udvardi, M. K., Czechowski, T., & Scheible, W.-R. (2008). Eleven golden rules of quantitative RT-PCR. *The Plant Cell*, 20(7), 1736–1737. DOI: [10.1105/tpc.108.061143](https://doi.org/10.1105/tpc.108.061143)
 34. Jain, M., Nijhawan, A., Tyagi, A. K., & Khurana, J. P. (2006). Validation of housekeeping genes as internal control for studying gene expression in rice by quantitative real-time PCR. *Biochemical and Biophysical Research Communications*, 345(2), 646–651. DOI: [10.1016/j.bbrc.2006.04.140](https://doi.org/10.1016/j.bbrc.2006.04.140)
 35. Kim, B.-R., Nam, H.-Y., Kim, S.-U., Kim, S.-I., & Chang, Y.-J. (2003). Normalization of reverse transcription quantitative-PCR with housekeeping genes in rice. *Biotechnology Letters*, 25(21), 1869–1872. DOI: [10.1023/A:1026298032009](https://doi.org/10.1023/A:1026298032009)
 36. Narsai, R., Ivanova, A., Ng, S., & Whelan, J. (2010). Defining reference genes in *Oryza sativa* using organ, development, biotic and abiotic transcriptome datasets. *BMC Plant Biology*, 10, 56. DOI: [10.1186/1471-2229-10-56](https://doi.org/10.1186/1471-2229-10-56)
 37. Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K., & Scheible, W.-R. (2005). Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiology*, 139(1), 5–17. DOI: [10.1104/pp.105.063743](https://doi.org/10.1104/pp.105.063743)
 38. Nicot, N., Hausman, J.-F., Hoffmann, L., & Evers, D. (2005). Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. *Journal of Experimental Botany*, 56(421), 2907–2914. DOI: [10.1093/jxb/eri285](https://doi.org/10.1093/jxb/eri285)
 39. Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., & Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology*, 3(7), research0034. DOI: [10.1186/gb-2002-3-7-research0034](https://doi.org/10.1186/gb-2002-3-7-research0034)
 40. Andersen, C. L., Jensen, J. L., & Ørntoft, T. F. (2004). Normalization of real-time quantitative reverse transcription-PCR data: A model-based variance estimation approach to identify genes suited for normalization. *Cancer Research*, 64(15), 5245–5250. DOI: [10.1158/0008-5472.CAN-04-0496](https://doi.org/10.1158/0008-5472.CAN-04-0496)
 41. Pfaffl, M. W., Tichopad, A., Prgomet, C., & Neuvians, T. P. (2004). Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper – Excel-based tool using pair-wise correlations. *Biotechnology Letters*, 26(6), 509–515. DOI: [10.1023/B:BILE.0000019559.84305.47](https://doi.org/10.1023/B:BILE.0000019559.84305.47)
 42. Hellemans, J., Mortier, G., De Paepe, A., Speleman, F., & Vandesompele, J. (2007). qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. *Genome Biology*, 8(2), R19. DOI: [10.1186/gb-2007-8-2-r19](https://doi.org/10.1186/gb-2007-8-2-r19)
 43. Bustin, S. A., Benes, V., Garson, J. A., Hellemans, J., Huggett, J., Kubista, M., ... Wittwer, C. T. (2009). The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry*, 55(4), 611–622. DOI: [10.1373/clinchem.2008.112797](https://doi.org/10.1373/clinchem.2008.112797)
 44. Bustin, S. A. (2010). Why the need for qPCR publication guidelines? The case for MIQE. *Methods*, 50(4), 217–226. DOI: [10.1016/j.ymeth.2009.12.006](https://doi.org/10.1016/j.ymeth.2009.12.006)