



ISSN : 2347 - 2243

Indo - American Journal of Life Sciences and Biotechnology



<https://iajlb.org>

Email : editor@iajlb.com or iajlb.editor@gmail.com



Algal Biofuels: Strain Selection, Cultivation Systems and Lipid Extraction

Elodie Boissonnet¹●, Marta Montalban²●, Elena Zanetti³●

¹Professor, Department of Industrial Microbiology, Vosges Institute of Applied Microbiology, Epinal, France.

²Senior Scientist, Department of Biotechnology, Rioja Institute of Biotechnology, Logrono, Spain.

³Scientist, Department of Industrial Microbiology, Monti Lepini Institute of Life Sciences, Frosinone, Italy.

Email: ¹e.boissonnet@viam-e.fr, ²m.montalban@ribt.es, ³e.zanetti@mlils.it

ORCID: ¹0000-0001-4431-8430, ²0000-0001-3097-7830, ³0000-0001-4136-2751

Received: February 26, 2019 Accepted: June 18, 2019 Published: October 14, 2019

DOI: [10.62644/iajlb.2019.v16.i4.pp11-18](https://doi.org/10.62644/iajlb.2019.v16.i4.pp11-18)

ABSTRACT

Microalgae are attractive as a source of liquid fuel because some strains accumulate large amounts of neutral lipid, grow on land that is unsuitable for crops and can use waste nutrients and carbon dioxide. Yet a decade of enthusiasm has not produced a commercial fuel that competes on price. This narrative review follows the chain from strain choice, through cultivation in open ponds and closed photobioreactors, to harvesting and lipid extraction, and asks where the real constraints lie. It draws on published work on algal biology, bioprocess engineering, extraction chemistry, techno-economic analysis and life-cycle assessment, and it reports no new data. Lipid content measured in a flask under nutrient stress is a poor guide to lipid productivity outdoors, because the stress that raises lipid content also slows growth. Open ponds are cheap but vulnerable to contamination, whereas closed systems are controllable but costly. Dewatering and extraction contribute a large share of energy use and cost, and no single solvent or disruption method suits every cell wall. The review concludes that algal biofuels remain technically plausible and economically unproven, that coproducts and waste-stream integration offer the more realistic route, and that results should be reported on a productivity basis with the whole process counted.

Keywords: Microalgae; Biofuel; Strain selection; Photobioreactor; Open pond; Lipid extraction; Narrative review

Citation: Boissonnet, Montalban and Zanetti [2019]. Algal Biofuels: Strain Selection, Cultivation Systems and Lipid Extraction. *Indo-Am. J. Life Sci. Biotechnol.* 16(4):11-18. DOI: [10.62644/iajlb.2019.v16.i4.pp11-18](https://doi.org/10.62644/iajlb.2019.v16.i4.pp11-18)

Article Class: Review Article

1. Introduction

Liquid fuel is hard to replace. Electricity can move a car, but ships, aircraft and heavy freight still depend on energy-dense liquids, and that fact has kept interest in biological fuels alive long after the first wave of enthusiasm for crop-based biodiesel faded (1, 2). Microalgae look like

an answer to the main objections. Some strains store a large share of their dry weight as lipid, they can grow in saline or wastewater on land that grows no food, and their photosynthetic efficiency is often described as higher than that of terrestrial crops (3, 4). Their lipids can be converted to the same kinds of fuel that are already burned in diesel engines (5).



The difficulty is the distance between a promising culture and a barrel of fuel. A strain that accumulates lipid happily in a flask may collapse in an outdoor pond, and a reactor that protects the culture may consume more energy than the oil is worth (6, 7). Reviews of the last decade agree that the field has been long on potential and short on demonstrated economics (8, 9).

This article is a narrative review of that chain, from the biology of lipid-rich algae to the chemistry of getting the oil out. It does not report new experiments, and it does not claim a systematic search. Where published work disagrees, we say so, and where the evidence depends on assumptions that are seldom stated, we try to name them. The aim is a fair account of what is known, not a catalogue of record claims.

2. Literature Review

2.1 Why Algae, and Which Algae

Only a small fraction of known algae have been examined as fuel feedstocks. The lipids of interest are mainly triacylglycerols, which many algae store when growth is limited by a nutrient such as nitrogen while light and carbon remain available (10, 11); their formation has been followed in detail in the model green alga *Chlamydomonas* (12). Membrane lipids and pigments come along in the extract but are a nuisance.

The fatty acid profile decides what the fuel will be like: chain length and unsaturation influence cold-flow behaviour, oxidative stability and ignition quality (13). Table 1 lists the traits that most often decide whether a strain is a realistic candidate.

2.2 Cultivation Systems

Two families of system dominate the literature. Open ponds, usually shallow

raceways stirred by a paddle wheel, are the older and cheaper design, and they have long been used for hardy species that thrive in extreme conditions (18, 19). Closed photobioreactors, whether tubular, flat-plate or column designs, protect the culture from invaders and allow better control of temperature and gas exchange, but they cost more to build and to run, and they bring their own problems of oxygen build-up and fouling (20, 21).

Table 1. Traits that make a microalgal strain a candidate for fuel production.

Trait	Why it matters	Common difficulty
Lipid productivity	Combines growth and lipid content (14)	High lipid often means slow growth
Robust growth outdoors	Must tolerate temperature and light swings (15, 16)	Flask winners may fail outdoors
Fatty acid profile	Sets fuel quality (13)	Too unsaturated for stability
Harvestability	Cell size and surface affect recovery (17)	Small cells stay suspended
Contamination resistance	Dominance is hard to hold in open systems (18)	Extreme conditions limit the choice

Note: Qualitative summary of the cited literature; no values are given because they depend on strain and conditions.

Neither system is simply better. A pond tolerates the weather and loses water by evaporation; a reactor excludes most contaminants and can overheat in sunshine. Such differences look small at flask scale but compound at the scale of a plant.

Heterotrophic growth in fermenters, where algae are fed an organic carbon source in the dark, avoids the light problem altogether at the price of a sugar bill (22). Table 2 sets the two main systems side by side.



Table 2. Open ponds and closed photobioreactors compared qualitatively.

Attribute	Open pond	Closed reactor
Capital cost	Lower (18, 21)	Higher (20)
Contamination	Frequent risk	Easier to exclude
Temperature control	Little; follows weather	Possible, at an energy cost
Gas transfer	Often limiting	Better but can build up oxygen
Scale-up	Straightforward in principle	Harder; many units or long tubes
Water loss	Evaporation can be large	Smaller

Note: Reading of the cited reviews; actual performance depends on site, strain and management.

2.3 Nutrients, Carbon and Wastewater

Algae need nitrogen, phosphorus and carbon in quantity, and supplying them from fertiliser and purchased carbon dioxide would count heavily against the fuel in any energy or cost balance (23, 24). Much recent work therefore proposes growing algae on wastewater, which supplies nutrients and treats the water at the same time (25, 26), with flue gas as a source of carbon dioxide that some strains answer with changed lipid accumulation (27).

The promise comes with qualifications. Wastewater varies from day to day, it carries its own microbial community, and a culture that must also meet discharge standards is not optimised for lipid. The question changes from how to produce oil to how to run a treatment process that yields oil as a byproduct. These points apply with equal force to any emerging bioenergy route, not only to algae.

3. Materials and Methods

3.1 Scope

The review covers microalgae grown for lipid-based fuels, mainly biodiesel and its

precursors: strain selection, cultivation, dewatering and lipid recovery, with techno-economic and life-cycle studies where they bear on those steps. Macroalgae, anaerobic digestion and thermochemical conversion lie outside the scope.

3.2 Literature Assembly

The sources are peer-reviewed reviews and research papers that the authors knew or located through citation chains, published up to the year of this issue. No formal systematic search was performed, so the selection reflects judgement and is open to the usual biases of a narrative review. None of these steps is difficult, but each is easy to skip.

3.3 Appraisal Framework

Each source was read against four questions: what was measured, at what scale, under what conditions, and how the result was expressed. A lipid percentage from a stressed flask culture, for instance, answers a different question from a productivity figure for an outdoor reactor over a season. Findings are presented qualitatively and grouped by step of the production chain; no numerical pooling was attempted. It is a habit worth keeping even when the answer seems obvious.

4. Results

4.1 Strain Selection and the Lipid Trade-off

The most robust finding in the strain literature is a trade-off. Conditions that push lipid content up, such as nitrogen starvation, tend to slow cell division, so oil accumulates in fewer cells. Lipid productivity, which combines growth rate and lipid content, is therefore the measure that matters (14), and screening programmes rank isolates on it under common conditions before testing the best



outdoors (15, 16). A culture that has had time to settle behaves differently from one that has not.

Even so, screening is fragile: a strain chosen in a controlled laboratory may rank differently when temperature, light and the resident microbial community change. Genetic engineering offers a way to separate growth from lipid accumulation (28, 29), and approaches that raise lipid by manipulating growth conditions are also reviewed (30). Engineered strains must still be shown to work outdoors, and releasing them into open ponds raises an unresolved regulatory question. Figure 1 places strain choice within the whole chain. Read this way, a lipid figure from a flask is a starting point, not a forecast.




Figure 1. Chain from strain to fuel and the places where losses and costs arise (schematic, not data).

4.2 Cultivation Performance

Reports from outdoor culture consistently show productivity falling short of laboratory expectation. Light is absorbed in the top layer of a dense culture, temperature swings with the day and the season, and grazers or competing algae can take over a pond within days (18, 20). Closed reactors improve control, yet the energy for mixing and cooling can offset

the gain, so net energy return depends heavily on design (23, 24). The point is not that a study is poor, only that its result travels less far than its abstract suggests.

Figure 2 summarises the qualitative picture. No cultivation mode is free of a concern; each mode trades one concern for another, and the choice depends on strain, climate and what else the plant is meant to do. Comparisons across studies therefore need the conditions stated. This is why the same species can look excellent in one report and unremarkable in the next.

4.3 Harvesting and Dewatering

Algal cultures are dilute, and the cells are small and slightly negatively charged, so they stay suspended. Recovering biomass is a major bottleneck, and flocculation, flotation, centrifugation and filtration each carry costs in energy, chemicals or contamination of the biomass (17, 31). Centrifugation is effective but energy-intensive; flocculation is cheaper but can leave residues that complicate later steps.

	Open pond	Closed reactor	Fermenter
Capital cost	low	high	high
Contamination risk	high	low	low
Needs light	yes	yes	no
Needs organic carbon	no	no	yes
Weather exposure	high	some	none

shading marks relative concern; schematic, not data

Figure 2. Qualitative matrix of cultivation modes against common concerns (schematic, not data).

Dewatering sits between cultivation and extraction, so its choice constrains both. Drying the paste before extraction is costly enough that many analyses treat it as decisive (32, 33), and no single harvesting



method is treated as solved.

4.4 Lipid Extraction


Lipid recovery from algae borrows from analytical chemistry. The chloroform and methanol procedures developed for animal tissue remain the reference against which other methods are compared (34, 35), but toxic solvents and low throughput make them unsuitable for a plant. Comparative studies find that the efficiency of a method depends on the species, on the state of the cells and on whether the biomass is wet or dry, and that no method wins everywhere (36, 37).

Cell wall strength is the main variable. Tough walls resist solvents, so mechanical disruption by bead milling, homogenisation, sonication or microwaves is often added, at the cost of energy (38, 39). Supercritical carbon dioxide is selective and leaves no solvent residue, but the pressure it needs raises capital cost (40). Table 3 sets out the main approaches, and Figure 3 gives a simple decision path.

Table 3. Lipid extraction approaches and the issues that accompany them.

Approach	Strength	Limitation
Chloroform–methanol	Thorough; the analytical reference (34, 35)	Toxic solvents; not for scale
Hexane after disruption	Less hazardous; suits fuel lipids (37)	Wet biomass extracts poorly
Mechanical disruption	Opens tough walls (36, 38)	Energy demand; heat
Supercritical carbon dioxide	Clean, selective (40)	High pressure and cost
Direct conversion of wet biomass	Skips drying (39)	Less mature

Note: Summary of reviews cited in the text; no performance figures are given.



schematic, not data

Figure 3. Decision tree for choosing a lipid recovery route (schematic, not data).

5. Discussion

5.1 Interpretation

Read as a whole, the literature supports a conditional reading. Algal lipids are real, and the biology is interesting, but the economics rest on assumptions that have rarely been tested at scale. Techno-economic studies reach different conclusions depending on the productivity assumed, the value of coproducts and the treatment of dewatering and extraction (32, 41). Life-cycle assessments find that the answer depends on how nutrients, drying and coproducts are handled, and that a poorly integrated system can use more fossil energy than it displaces (23, 24). A saving that vanishes once the whole process is counted is not one an operator would ever see.

The more realistic route appears to be integration: producing fuel alongside higher-value products, or treating water while producing oil, so that fuel is one revenue among several (19, 42). Several authors suggest that fuel may follow only after a biorefinery model is established (7, 43), and fuel from algal oil must in any case meet quality standards that raw oil often does not (5, 44). Nothing here argues



against algal fuels; it argues for measuring what happens at scale.

5.2 Limitations

This review has the limitations of its type. Sources were selected without a systematic search, so relevant work may be missing. Many studies are laboratory or pilot studies using different strains, units and conditions, which prevents direct comparison, and economic and life-cycle studies depend on modelling assumptions that vary widely. We have avoided quoting numbers whose provenance we could not confirm, which makes the account less specific than a reader might wish, but safer. Reviews are more useful when they say where the evidence stops than when they smooth it over.

5.3 Future Work

Three needs stand out. First, shared reporting standards, so that studies state productivity per unit area and time and describe the conditions (45, 46). Second, long outdoor trials with realistic contamination. Third, integrated assessments that count dewatering, extraction, nutrients and coproducts in a single balance (43, 47).

6. Conclusion

Microalgae can make lipid suitable for fuel, and several strains and systems have been shown to work in demonstration. What the literature has not shown is a route that is competitive without subsidy or coproduct revenue. The central difficulty is not that algae cannot make oil, but that the conditions for high lipid content, the cost of cultivation, and the energy for dewatering and extraction pull against each other. Researchers should report productivity, state conditions and count the whole process; planners should treat algal fuel as an option tied to waste treatment or coproducts. That is a modest conclusion,

and it is the one the literature will bear.

7. Acknowledgements

The authors thank colleagues who commented on earlier drafts of this review.

8. Funding

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

9. Conflict of Interest

The authors declare no conflict of interest.

10. Author Contributions

E.B. conceived the review and wrote the sections on strain selection and cultivation. M.M. wrote the sections on harvesting and process integration. E.Z. wrote the sections on lipid extraction and appraisal. All authors revised the manuscript and approved the final version.

11. Ethics Statement

This is a review of published literature. No human participants or animals were involved and no ethical approval was required.

12. Data Availability

No new data were generated. All sources are cited in the reference list.

13. References

1. Chisti, Y. (2007). Biodiesel from microalgae. *Biotechnology Advances*, 25(3), 294–306. DOI: [10.1016/j.biotechadv.2007.02.001](https://doi.org/10.1016/j.biotechadv.2007.02.001)
2. Chisti, Y. (2008). Biodiesel from microalgae beats bioethanol. *Trends in Biotechnology*, 26(3), 126–131. DOI: [10.1016/j.tibtech.2007.12.002](https://doi.org/10.1016/j.tibtech.2007.12.002)
3. Mata, T. M., Martins, A. A., & Caetano, N. S. (2010). Microalgae for biodiesel production and other applications: A review. *Renewable and Sustainable Energy Reviews*, 14(1), 217–232. DOI: [10.1016/j.rser.2009.07.020](https://doi.org/10.1016/j.rser.2009.07.020)
4. Brennan, L., & Owende, P. (2010). Biofuels from microalgae: A review of technologies for production, processing, and extractions of biofuels and co-products. *Renewable and Sustainable Energy Reviews*, 14(2), 557–577. DOI: [10.1016/j.rser.2009.10.009](https://doi.org/10.1016/j.rser.2009.10.009)



5. Gouveia, L., & Oliveira, A. C. (2009). Microalgae as a raw material for biofuels production. *Journal of Industrial Microbiology and Biotechnology*, 36(2), 269–274. DOI: [10.1007/s10295-008-0495-6](https://doi.org/10.1007/s10295-008-0495-6)
6. Wijffels, R. H., & Barbosa, M. J. (2010). An outlook on microalgal biofuels. *Science*, 329(5993), 796–799. DOI: [10.1126/science.1189003](https://doi.org/10.1126/science.1189003)
7. Chisti, Y. (2013). Constraints to commercialization of algal fuels. *Journal of Biotechnology*, 167(3), 201–214. DOI: [10.1016/j.jbiotec.2013.07.020](https://doi.org/10.1016/j.jbiotec.2013.07.020)
8. Pienkos, P. T., & Darzins, A. (2009). The promise and challenges of microalgal-derived biofuels. *Biofuels, Bioproducts and Biorefining*, 3(4), 431–440. DOI: [10.1002/bbb.159](https://doi.org/10.1002/bbb.159)
9. Hannon, M., Gimpel, J., Tran, M., Rasala, B., & Mayfield, S. (2010). Biofuels from algae: Challenges and potential. *Biofuels*, 1(5), 763–784. DOI: [10.4155/bfs.10.44](https://doi.org/10.4155/bfs.10.44)
10. Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M., & Darzins, A. (2008). Microalgal triacylglycerols as feedstocks for biofuel production: Perspectives and advances. *The Plant Journal*, 54(4), 621–639. DOI: [10.1111/j.1365-3113X.2008.03492.x](https://doi.org/10.1111/j.1365-3113X.2008.03492.x)
11. Guschina, I. A., & Harwood, J. L. (2006). Lipids and lipid metabolism in eukaryotic algae. *Progress in Lipid Research*, 45(2), 160–186. DOI: [10.1016/j.plipres.2006.01.001](https://doi.org/10.1016/j.plipres.2006.01.001)
12. Wang, Z. T., Ullrich, N., Joo, S., Waffenschmidt, S., & Goodenough, U. (2009). Algal lipid bodies: Stress induction, purification, and biochemical characterization in wild-type and starchless *Chlamydomonas reinhardtii*. *Eukaryotic Cell*, 8(12), 1856–1868. DOI: [10.1128/EC.00272-09](https://doi.org/10.1128/EC.00272-09)
13. Knothe, G. (2005). Dependence of biodiesel fuel properties on the structure of fatty acid alkyl esters. *Fuel Processing Technology*, 86(10), 1059–1070. DOI: [10.1016/j.fuproc.2004.11.002](https://doi.org/10.1016/j.fuproc.2004.11.002)
14. Griffiths, M. J., & Harrison, S. T. L. (2009). Lipid productivity as a key characteristic for choosing algal species for biodiesel production. *Journal of Applied Phycology*, 21(5), 493–507. DOI: [10.1007/s10811-008-9392-7](https://doi.org/10.1007/s10811-008-9392-7)
15. Rodolfi, L., Chini Zittelli, G., Bassi, N., Padovani, G., Biondi, N., Bonini, G., & Tredici, M. R. (2009). Microalgae for oil: Strain selection, induction of lipid synthesis and outdoor mass cultivation in a low-cost photobioreactor. *Biotechnology and Bioengineering*, 102(1), 100–112. DOI: [10.1002/bit.22033](https://doi.org/10.1002/bit.22033)
16. Mutanda, T., Ramesh, D., Karthikeyan, S., Kumari, S., Anandraj, A., & Bux, F. (2011). Bioprospecting for hyper-lipid producing microalgal strains for sustainable biofuel production. *Bioresource Technology*, 102(1), 57–70. DOI: [10.1016/j.biortech.2010.06.077](https://doi.org/10.1016/j.biortech.2010.06.077)
17. Christenson, L., & Sims, R. (2011). Production and harvesting of microorganisms for biofuel production. *Biotechnology Advances*, 29(6), 686–702. DOI: [10.1016/j.biotechadv.2011.05.015](https://doi.org/10.1016/j.biotechadv.2011.05.015)
18. Borowitzka, M. A. (1999). Commercial production of microalgae: Ponds, tanks, tubes and fermenters. *Journal of Biotechnology*, 70(1–3), 313–321. DOI: [10.1016/S0168-1656\(99\)00083-8](https://doi.org/10.1016/S0168-1656(99)00083-8)
19. Borowitzka, M. A. (2013). High-value products from microalgae: Their development and commercialisation. *Journal of Applied Phycology*, 25(3), 743–756. DOI: [10.1007/s10811-013-9983-9](https://doi.org/10.1007/s10811-013-9983-9)
20. Pulz, O. (2001). Photobioreactors: Production systems for phototrophic microorganisms. *Applied Microbiology and Biotechnology*, 57(3), 287–293. DOI: [10.1007/s002530100702](https://doi.org/10.1007/s002530100702)
21. Ugwu, C. U., Aoyagi, H., & Uchiyama, H. (2008). Photobioreactors for mass cultivation of algae. *Bioresource Technology*, 99(10), 4021–4028. DOI: [10.1016/j.biortech.2007.01.046](https://doi.org/10.1016/j.biortech.2007.01.046)
22. Xu, H., Miao, X., & Wu, Q. (2006). High quality biodiesel production from a microalga *Chlorella protothecoides* by heterotrophic growth in fermenters. *Journal of Biotechnology*, 126(4), 499–507. DOI: [10.1016/j.jbiotec.2006.05.002](https://doi.org/10.1016/j.jbiotec.2006.05.002)
23. Lardon, L., Hélias, A., Sialve, B., Steyer, J.-P., & Bernard, O. (2009). Life-cycle assessment of biodiesel production from microalgae. *Environmental Science & Technology*, 43(17), 6475–6481. DOI: [10.1021/es900705j](https://doi.org/10.1021/es900705j)
24. Clarens, A. F., Resurreccion, E. P., White, M. A., & Colosi, L. M. (2010). Environmental life cycle comparison of algae to other bioenergy feedstocks. *Environmental Science & Technology*, 44(5), 1813–1819. DOI: [10.1021/es902838n](https://doi.org/10.1021/es902838n)
25. Pittman, J. K., Dean, A. P., & Osundeko, O. (2011). The potential of sustainable algal biofuel production using wastewater resources. *Bioresource Technology*, 102(1), 17–25. DOI: [10.1016/j.biortech.2010.06.035](https://doi.org/10.1016/j.biortech.2010.06.035)
26. Rawat, I., Ranjith Kumar, R., Mutanda, T., & Bux, F. (2011). Dual role of microalgae: Phycoremediation of domestic wastewater and biomass production for sustainable biofuels production. *Applied Energy*, 88(10), 3411–3424. DOI: [10.1016/j.apenergy.2010.11.025](https://doi.org/10.1016/j.apenergy.2010.11.025)
27. Chiu, S.-Y., Kao, C.-Y., Tsai, M.-T., Ong, S.-C., Chen, C.-H., & Lin, C.-S. (2009). Lipid accumulation and CO₂ utilization of *Nannochloropsis oculata* in response to CO₂ aeration. *Bioresource Technology*, 100(2), 833–838. DOI: [10.1016/j.biortech.2008.06.061](https://doi.org/10.1016/j.biortech.2008.06.061)



28. Radakovits, R., Jinkerson, R. E., Darzins, A., & Posewitz, M. C. (2010). Genetic engineering of algae for enhanced biofuel production. *Eukaryotic Cell*, 9(4), 486–501. DOI: [10.1128/EC.00364-09](https://doi.org/10.1128/EC.00364-09)
29. Georgianna, D. R., & Mayfield, S. P. (2012). Exploiting diversity and synthetic biology for the production of algal biofuels. *Nature*, 488(7411), 329–335. DOI: [10.1038/nature11479](https://doi.org/10.1038/nature11479)
30. Sharma, K. K., Schuhmann, H., & Schenk, P. M. (2012). High lipid induction in microalgae for biodiesel production. *Energies*, 5(5), 1532–1553. DOI: [10.3390/en5051532](https://doi.org/10.3390/en5051532)
31. Uduman, N., Qi, Y., Danquah, M. K., Forde, G. M., & Hoadley, A. (2010). Dewatering of microalgal cultures: A major bottleneck to algae-based fuels. *Journal of Renewable and Sustainable Energy*, 2(1), 012701. DOI: [10.1063/1.3294480](https://doi.org/10.1063/1.3294480)
32. Davis, R., Aden, A., & Pienkos, P. T. (2011). Techno-economic analysis of autotrophic microalgae for fuel production. *Applied Energy*, 88(10), 3524–3531. DOI: [10.1016/j.apenergy.2011.04.018](https://doi.org/10.1016/j.apenergy.2011.04.018)
33. Williams, P. J. I. B., & Laurens, L. M. L. (2010). Microalgae as biodiesel and biomass feedstocks: Review and analysis of the biochemistry, energetics and economics. *Energy & Environmental Science*, 3(5), 554–590. DOI: [10.1039/B924978H](https://doi.org/10.1039/B924978H)
34. Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8), 911–917. DOI: [10.1139/o59-099](https://doi.org/10.1139/o59-099)
35. Folch, J., Lees, M., & Sloane Stanley, G. H. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *Journal of Biological Chemistry*, 226(1), 497–509. DOI: [10.1016/S0021-9258\(18\)64849-5](https://doi.org/10.1016/S0021-9258(18)64849-5)
36. Lee, J.-Y., Yoo, C., Jun, S.-Y., Ahn, C.-Y., & Oh, H.-M. (2010). Comparison of several methods for effective lipid extraction from microalgae. *Bioresource Technology*, 101(1), S75–S77. DOI: [10.1016/j.biortech.2009.03.058](https://doi.org/10.1016/j.biortech.2009.03.058)
37. Halim, R., Danquah, M. K., & Webley, P. A. (2012). Extraction of oil from microalgae for biodiesel production: A review. *Biotechnology Advances*, 30(3), 709–732. DOI: [10.1016/j.biotechadv.2012.01.001](https://doi.org/10.1016/j.biotechadv.2012.01.001)
38. Mercer, P., & Armenta, R. E. (2011). Developments in oil extraction from microalgae. *European Journal of Lipid Science and Technology*, 113(5), 539–547. DOI: [10.1002/ejlt.201000455](https://doi.org/10.1002/ejlt.201000455)
39. Ranjith Kumar, R., Hanumantha Rao, P., & Arumugam, M. (2015). Lipid extraction methods from microalgae: A comprehensive review. *Frontiers in Energy Research*, 2, 61. DOI: [10.3389/fenrg.2014.00061](https://doi.org/10.3389/fenrg.2014.00061)
40. Cooney, M., Young, G., & Nagle, N. (2009). Extraction of bio-oils from microalgae. *Separation and Purification Reviews*, 38(4), 291–325. DOI: [10.1080/15422110903327919](https://doi.org/10.1080/15422110903327919)
41. Stephens, E., Ross, I. L., King, Z., Mussnug, J. H., Kruse, O., Posten, C., ... Hankamer, B. (2010). An economic and technical evaluation of microalgal biofuels. *Nature Biotechnology*, 28(2), 126–128. DOI: [10.1038/nbt0210-126](https://doi.org/10.1038/nbt0210-126)
42. Spolaore, P., Joannis-Cassan, C., Duran, E., & Isambert, A. (2006). Commercial applications of microalgae. *Journal of Bioscience and Bioengineering*, 101(2), 87–96. DOI: [10.1263/jbb.101.87](https://doi.org/10.1263/jbb.101.87)
43. Slade, R., & Bauen, A. (2013). Micro-algae cultivation for biofuels: Cost, energy balance, environmental impacts and future prospects. *Biomass and Bioenergy*, 53, 29–38. DOI: [10.1016/j.biombioe.2012.12.019](https://doi.org/10.1016/j.biombioe.2012.12.019)
44. Li, Y., Horsman, M., Wu, N., Lan, C. Q., & Dubois-Calero, N. (2008). Biofuels from microalgae. *Biotechnology Progress*, 24(4), 815–820. DOI: [10.1021/bp070371k](https://doi.org/10.1021/bp070371k)
45. Lam, M. K., & Lee, K. T. (2012). Microalgae biofuels: A critical review of issues, problems and the way forward. *Biotechnology Advances*, 30(3), 673–690. DOI: [10.1016/j.biotechadv.2011.11.008](https://doi.org/10.1016/j.biotechadv.2011.11.008)
46. Scott, S. A., Davey, M. P., Dennis, J. S., Horst, I., Howe, C. J., Lea-Smith, D. J., & Smith, A. G. (2010). Biodiesel from algae: Challenges and prospects. *Current Opinion in Biotechnology*, 21(3), 277–286. DOI: [10.1016/j.copbio.2010.03.005](https://doi.org/10.1016/j.copbio.2010.03.005)
47. Singh, J., & Gu, S. (2010). Commercialization potential of microalgae for biofuels production. *Renewable and Sustainable Energy Reviews*, 14(9), 2596–2610. DOI: [10.1016/j.rser.2010.06.014](https://doi.org/10.1016/j.rser.2010.06.014)